(a)
$$m = \frac{W}{g} = \frac{3500}{32.2} = \frac{108.7 \text{ slugs}}{108.7 \text{ slugs}}$$

(b) $W = 3500 \text{ lb} \left[\frac{4.4482 \text{ N}}{1 \text{ lb}} \right] = \frac{15570 \text{ N}}{9.81}$
(c) $m = \frac{W}{g} = \frac{15570}{9.81} = \frac{1587 \text{ kg}}{9.81}$

For a
$$180-16$$
 person:
 $W = mg$: $180 \cdot 16 = m (32.2 \text{ ft/sec}^2)$
 $m = 5.59 \text{ slugs}$
 $180 \cdot 16 \left(\frac{4.4482 \text{ N}}{16}\right) = \frac{801 \text{ N}}{16}$
 $W = mg$: $801 \cdot N = m (9.81 \text{ m/s}^2)$
 $m = 81.6 \text{ kg}$

$$\frac{1/3}{V_1} = \frac{15}{5} \left(\frac{4}{5} + \frac{3}{5} \right) = \frac{12}{5} + \frac{9}{1}$$

$$\frac{V_2}{V_2} = \frac{12}{5} \left(-\cos 60^{\circ} + \sin 60^{\circ} \right) = -6 + 10.39 = -6 + 10.$$

The weight of an average apple is
$$W = \frac{5 \text{ lb}}{12 \text{ apples}} = 0.417 \text{ lb}$$

Mass in slugs is $m = \frac{W}{g} = \frac{0.417}{32.2} = 0.01294 \text{ slugs}$

Mass in kg is $m = 0.01294 \text{ slugs} \left(\frac{14.594 \text{ kg}}{1 \text{ slug}}\right)$
 $= 0.1888 \text{ kg}$

Weight in N is $W = mg = 0.1888 \left(9.81\right) = 1.853 \text{ N}$

These apples weigh closer to 2 N each than to the rule of 1 N each!

1/5 Mass of iron sphere
$$m = PV$$

= $(7210 \frac{kg}{m^3}) (\frac{4}{3} \% (0.050)^3) = 3.78 \text{ kg}$
Force of mutual attraction: $\frac{Gm^2}{d^2}$
Weight of each sphere: $\frac{Gmem}{r^2}$
 $\frac{Gm^2}{d^2} = \frac{Gmem}{r^2}$, $r = d\sqrt{\frac{me}{m}}$
= $0.1\sqrt{\frac{5.976 \times 10^{24}}{3.78}} \frac{1}{10^3}$
= $1.258 (10^8) \text{ km}$

$$\frac{1/6}{6R}$$

$$\frac{35^{\circ}}{8} - - \times \\
F = \frac{6m_{T_{i}}m_{Cu}}{d^{2}} = \frac{6\left[\frac{4}{3}\pi R^{3}\Gamma_{T_{i}}\right]\left[\frac{4}{3}\pi (2R)^{3}\Gamma_{Cu}\right]}{(6R)^{2}}$$

$$= \frac{32}{81}\pi^{2}G\Gamma_{T_{i}}\Gamma_{Cu}R^{4}$$

$$= \frac{32}{81}\pi^{2}(6.673\cdot10^{-11})(4510)(8910)(0.040)^{4}$$

$$= 2.68(10^{-8})N$$
Force is a vector quantity, so
$$F = F_{D} = 2.68(10^{-8})\left[-\cos 35^{\circ}i - \sin 35^{\circ}i\right]$$

$$= (-2.19i - 1.535j)10^{-8}N$$

1/7
$$mg = \frac{1}{3}mg_{h=0}$$

 $\frac{R^2}{(R+h)^2}g_0 = \frac{1}{3}g_0$
Solve for $h: h = (-13-1)R = 0.732R$

$$g_{rel} = 9.780 \, 327 \, (1 + 0.005 \, 279 \, \sin^2 8 + 0.000 \, 023 \, \sin^4 8 \, \cdots)$$
 $A \pm 8 = 35^{\circ}$, $g_{rel} = 9.797 \, 337 \, m/s^2$
 $g_{abs} = g_{rel} + 0.03382 \, \cos^2 8$
 $= 9.797 \, 337 + 0.03382 \, \cos^2 35^{\circ}$
 $= 9.820 \, 031 \, m/s^2$
 $W_{abs} = mg_{abs} = 60 \, (9.820 \, 031) = 589 \, N$
 $W_{rel} = mg_{rel} = 60 \, (9.797 \, 337) = 588 \, N$

(More precise values: $W_{abs} = 589.2 \, N$
 $W_{rel} = 587.8 \, N$)

$$\frac{1/9}{9h} = \frac{Gme}{(R+h)^2}$$

$$= \frac{(3.439 \cdot 10^{-8})(4.095 \cdot 10^{23})}{[(3959 + 200)(5280)]^2} = 29.2 \text{ ft/sec}^2$$
Mass of passenger $m = \frac{W}{9} = \frac{180}{32.174}$

$$= 5.59 \text{ slugs}$$

Absolute weight at h = 200 mi: $W_h = mg_h = (5.59)(29.2) = 163.4 \text{ lb}$ The terms "zero-g" and "weightless"

are absolutely (!) misnomers in this case.

