# Fox and McDonalds Introduction to Fluid Mechanics 8th Edition Pritchard Solutions Manual

Full Download: http://alibabadownload.com/product/fox-and-mcdonalds-introduction-to-fluid-mechanics-8th-edition-pritchard-sol

# Problem 2.33

[Difficulty: 3]

<b>2.33</b> A flow is $a = 1/5 \text{ s}^{-1}$ and meters. Obtain through point the particle to $(1, 1)$ ? What streamline and ticle. What constreamline, and	described by nd $b=1$ m/ in the equa (1, 1). At $t=$ hat initially t are its co d the initial, onclusions c: d streakline	velocity field $\vec{V} = ax\hat{i} + i$ s. Coordinates are me tion for the streamlin = 5 s, what are the coor (at $t=0$ ) passed thro ordinates at $t=10$ s? 5 s, and 10 s positions of an you draw about the for this flow?	$b\hat{j}$ , where assured in the passing rdinates of ugh point Plot the pof the par- the pathline,				
Given:	Velocity fi	eld					
Find:	Equation for compare particular	Equation for streamline through point (1.1); coordinates of particle at $t = 5$ s and $t = 10$ s that was at (1,1) at $t = 0$ ; compare pathline, streamline, streakline					
Solution:							
Governing equ	ations:	For streamlines	$\frac{v}{u} = \frac{dy}{dx}$	For pathlines	$u_p = \frac{dx}{dt}$	v <sub>p</sub> =	dy dt
Assumption: 2	2D flow						
Given data		$a = \frac{1}{5} \frac{1}{s} \qquad b = 1$	$\frac{m}{s}$ $x_0 = 1$	$y_0 = 1$	$t_0 = 0$		
For streamlines		$\frac{v}{u} = \frac{dy}{dx} = \frac{b}{a \cdot x}$					
So, separating v	variables	$\frac{a}{b} \cdot dy = \frac{dx}{x}$					
Integrating		$\frac{a}{b} \cdot \left( y - y_0 \right) = \ln \left( \frac{x}{x_0} \right)$	) /				
The solution is	then	$y = y_0 + \frac{b}{a} \cdot \ln\left(\frac{x}{x_0}\right)$	$= 5 \cdot \ln(x) + 1$				
Hence for pathl	ines	$u_p = \frac{dx}{dt} = a \cdot x$		$v_p = \frac{dy}{dt} = b$			
Hence		$\frac{\mathrm{d}x}{x} = a \cdot \mathrm{d}t$		$dy = b \cdot dt$			
Integrating		$\ln\!\!\left(\frac{x}{x_0}\right) = a \cdot \! \left(t - t_0\right)$		$y - y_0 = b \cdot (t - t)$	$(t_0)$		
The pathlines a	re	$\mathbf{x} = \mathbf{x}_0 \cdot \mathbf{e}^{\mathbf{a} \cdot \left(t - t_0\right)}$		$y = y_0 + b \cdot (t - t)$	$(t_0)$	or	$y = y_0 + \frac{b}{a} \cdot \ln\left(\frac{x}{x_0}\right)$

# This sample only, Download all chapters at: alibabadownload.com

For a particle that was at  $x_0 = 1$  m,  $y_0 = 1$  m at  $t_0 = 0$  s, at time t = 1 s we find the position is

$$x = x_0 \cdot e^{a \cdot (t-t_0)} = e^{\frac{1}{5}} m$$
  $y = y_0 + b \cdot (t-t_0) = 2 m$ 

For a particle that was at  $x_0 = 1$  m,  $y_0 = 1$  m at  $t_0 = 0$  s, at time t = 5 s we find the position is

$$x = x_0 e^{a \cdot (t-t_0)} = e m \qquad y = y_0 + b \cdot (t-t_0) = 6 m$$

For a particle that was at  $x_0 = 1$  m,  $y_0 = 1$  at  $t_0 = 0$  s, at time t = 10 s we find the position is

$$x = x_0 \cdot e^{a \cdot (t-t_0)} = e^2 m$$
  $y = y_0 + b \cdot (t-t_0) = 11 m$ 

For this steady flow streamlines, streaklines and pathlines coincide



**Streamline and Position Plots** 

**2.34** A flow is described by velocity field  $\vec{V} = a\hat{i} + bx\hat{j}$ , where a = 2 m/s and b = 1 s<sup>-1</sup>. Coordinates are measured in meters. Obtain the equation for the streamline passing through point (2, 5). At t = 2 s, what are the coordinates of the particle that passed through point (0, 4) at t = 0? At t = 3 s, what are the coordinates of the particle that passed through point (1, 4.25) 2 s earlier? What conclusions can you draw about the pathline, streamline, and streakline for this flow?

**Given:** Velocity field

**Find:** Equation for streamline through point (2.5); coordinates of particle at t = 2 s that was at (0,4) at t = 0; coordinates of particle at t = 3 s that was at (1,4.25) at t = 1 s; compare pathline, streamline, streakline

# Solution:

Governing equations:	For streamlines $\frac{v}{u} = \frac{dy}{dx}$	For pathlines $u_p = \frac{dx}{dt}$ $v_p = \frac{dy}{dt}$
Assumption: 2D flow		
Given data	$a = 2 \frac{m}{s}$ $b = 1 \frac{1}{s}$ $x_0 = 2$	$y_0 = 5$ $x = 1$ $x = x$
For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}} = \frac{\mathbf{b}\cdot\mathbf{x}}{\mathbf{a}}$	
So, separating variables	$\frac{a}{b} \cdot dy = x \cdot dx$	
Integrating	$\frac{a}{b} \cdot \left( y - y_0 \right) = \frac{1}{2} \cdot \left( x^2 - x_0^2 \right)$	
The solution is then	$y = y_0 + \frac{b}{2 \cdot a} \cdot \left(x^2 - x_0^2\right) = \frac{x^2}{4} + \frac{b}{4}$	4
Hence for pathlines	$u_p = \frac{dx}{dt} = a$	$\mathbf{v}_{\mathbf{p}} = \frac{\mathrm{d}\mathbf{y}}{\mathrm{d}t} = \mathbf{b} \cdot \mathbf{x}$
Hence	$d\mathbf{x} = \mathbf{a} \cdot d\mathbf{t}$	$dy = b \cdot x \cdot dt$
Integrating	$x - x_0 = a \cdot \left(t - t_0\right)$	$dy = b \cdot \left[ x_0 + a \cdot \left( t - t_0 \right) \right] \cdot dt$
		$y - y_0 = b \cdot \left[ x_0 \cdot (t - t_0) + \frac{a}{2} \cdot \left( \left( t^2 - t_0^2 \right) \right) - a \cdot t_0 \cdot (t - t_0) \right]$
The pathlines are	$\mathbf{x} = \mathbf{x}_0 + \mathbf{a} \cdot \left( \mathbf{t} - \mathbf{t}_0 \right)$	$y = y_0 + b \cdot \left[ x_0 \cdot (t - t_0) + \frac{a}{2} \cdot \left( \left( t^2 - t_0^2 \right) \right) - a \cdot t_0 \cdot (t - t_0) \right]$

For a particle that was at  $x_0 = 0$  m,  $y_0 = 4$  m at  $t_0 = 0$ s, at time t = 2 s we find the position is

$$x = x_0 + a \cdot (t - t_0) = 4m \qquad y = y_0 + b \cdot \left[ x_0 \cdot (t - t_0) + \frac{a}{2} \cdot \left( \left( t^2 - t_0^2 \right) \right) - a \cdot t_0 \cdot (t - t_0) \right] = m$$

For a particle that was at  $x_0 = 1 \text{ m}$ ,  $y_0 = 4.25 \text{ m}$  at  $t_0 = 1 \text{ s}$ , at time t = 3 s we find the position is

$$x = x_0 + a \cdot (t - t_0) = 5 m \qquad y = y_0 + b \cdot \left[ x_0 \cdot (t - t_0) + \frac{a}{2} \cdot \left( \left( t^2 - t_0^2 \right) \right) - a \cdot t_0 \cdot (t - t_0) \right] = 10. m$$

For this steady flow streamlines, streaklines and pathlines coincide; the particles refered to are the same particle!



**2.35** A flow is described by velocity field  $\vec{V} = ay\hat{i} + bt\hat{j}$ , where  $a = 0.2 \text{ s}^{-1}$  and  $b = 0.4 \text{ m/s}^2$ . At t = 2 s, what are the coordinates of the particle that passed through point (1, 2) at t = 0? At t = 3 s, what are the coordinates of the particle that passed through point (1, 2) at t = 2 s? Plot the pathline and streakline through point (1, 2), and plot the streamlines through the same point at the instants t = 0, 1, 2, and 3 s.

**Given:** Velocity field

**Find:** Coordinates of particle at t = 2 s that was at (1,2) at t = 0; coordinates of particle at t = 3 s that was at (1,2) at t = 2 s; plot pathline and streakline through point (1,2) and compare with streamlines through same point at t = 0, 1 and 2 s

### Solution

Governing equations:	For pathlines	$u = \frac{dx}{dx}$	$v = \frac{dy}{dy}$	For	<u>v</u> =	dy
8 1	1	<sup>ap</sup> dt	<sup>p</sup> dt	streamlines	u	dx

Following the discussion leading up to Eq. 2.10, we first find equations for the pathlines in form

 $\mathbf{x}_{p}(t) = \mathbf{x} \Big( t, \mathbf{x}_{0}, \mathbf{y}_{0}, t_{0} \Big) \qquad \text{and} \qquad \qquad \mathbf{y}_{p}(t) = \mathbf{y} \Big( t, \mathbf{x}_{0}, \mathbf{y}_{0}, t_{0} \Big)$ 

 $x_{st}\!\left(t_0\right) = x\!\left(t, x_0, y_0, t_0\right) \quad \text{ and } \quad y_{st}\!\left(t_0\right) = y\!\left(t, x_0, y_0, t_0\right)$ 

which gives the streakline at t, where  $x_0$ ,  $y_0$  is the point at which dye is released ( $t_0$  is varied from 0 to t)

Assumption: 2D flow

Given data  $a = 0.2 \quad \frac{1}{s} \quad b = 0.4 \quad \frac{m}{s^{2}}$ Hence for pathlines  $u_{p} = \frac{dx}{dt} = a \cdot y \qquad v_{p} = \frac{dy}{dt} = b \cdot t$ Hence  $dx = a \cdot y \cdot dt \qquad dy = b \cdot t \cdot dt \qquad y - y_{0} = \frac{b}{2} \cdot \left(t^{2} - t_{0}^{2}\right)$ For x  $dx = \left[a \cdot y_{0} + a \cdot \frac{b}{2} \cdot \left(t^{2} - t_{0}^{2}\right)\right] \cdot dt$ Integrating  $x - x_{0} = a \cdot y_{0} \cdot \left(t - t_{0}\right) + a \cdot \frac{b}{2} \cdot \left[\frac{t^{3}}{3} - \frac{t_{0}^{3}}{3} - t_{0}^{2} \cdot \left(t - t_{0}\right)\right]$ The pathlines are  $x(t) = x_{0} + a \cdot y_{0} \cdot \left(t - t_{0}\right) + a \cdot \frac{b}{2} \cdot \left[\frac{t^{3}}{3} - \frac{t_{0}^{3}}{3} - t_{0}^{2} \cdot \left(t - t_{0}\right)\right]$   $y(t) = y_{0} + \frac{b}{2} \cdot \left(t^{2} - t_{0}^{2}\right)$ 

These give the position (x,y) at any time t of a particle that was at  $(x_0,y_0)$  at time  $t_0$ 

Note that streaklines are obtained using the logic of the Governing equations, above

The streak lines are 
$$x(t_0) = x_0 + a \cdot y_0 \cdot (t - t_0) + a \cdot \frac{b}{2} \cdot \left[ \frac{t^3}{3} - \frac{t_0^3}{3} - t_0^2 \cdot (t - t_0) \right]$$
  $y(t_0) = y_0 + \frac{b}{2} \cdot \left( t^2 - t_0^2 \right)$ 

These gives the streakline at t, where  $x_0$ ,  $y_0$  is the point at which dye is released ( $t_0$  is varied from 0 to t)

For a particle that was at  $x_0 = 1 \text{ m}$ ,  $y_0 = 2 \text{ m}$  at  $t_0 = 0$ s, at time t = 2 s we find the position is (from pathline equations)

$$x = x_0 + a \cdot y_0 \cdot (t - t_0) + a \cdot \frac{b}{2} \cdot \left[ \frac{t^3}{3} - \frac{t_0^3}{3} - t_0^2 \cdot (t - t_0) \right] = 1.9 \text{ m} \qquad y = y_0 + \frac{b}{2} \cdot (t^2 - t_0^2) = 2.8 \text{ m}$$

For a particle that was at  $x_0 = 1 \text{ m}$ ,  $y_0 = 2 \text{ m at } t_0 = 2 \text{ s}$ , at time t = 3 s we find the position is

$$x = x_0 + a \cdot y_0 \cdot (t - t_0) + a \cdot \frac{b}{2} \cdot \left[ \frac{t^3}{3} - \frac{t_0^3}{3} - t_0^2 \cdot (t - t_0) \right] = 1.4 \text{ m} \qquad y = y_0 + \frac{b}{2} \cdot (t^2 - t_0^2) = 3.0 \text{ m}$$

For streamlines

Integrating

$$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}} = \frac{\mathbf{b}\cdot\mathbf{t}}{\mathbf{a}\cdot\mathbf{y}}$$

So, separating variables  $y \cdot dy = \frac{b}{a} \cdot t \cdot dx$  where we treat t as a constant

$$\frac{y^2 - y_0^2}{2} = \frac{b \cdot t}{a} \cdot (x - x_0) \quad \text{and we have} \quad x_0 = 1 \text{ m } y_0 = 2 \text{ m}$$

The streamlines are then 
$$y = \sqrt{y_0^2 + \frac{2 \cdot b \cdot t}{a} \cdot (x - x_0)} = \sqrt{4 \cdot t \cdot (x - 1) + 4}$$





2.1 For the velocity fields given below, determine:

- a. whether the flow field is one-, two-, or three-dimensional, and why.
- b. whether the flow is steady or unsteady, and why. (The quantities a and b are constants.)

(1)  $\vec{V} = [(ax + t)e^{by}]\hat{i}$  (2)  $\vec{V} = (ax - by)\hat{i}$ (3)  $\vec{V} = ax\hat{i} + [e^{bx}]\hat{j}$  (4)  $\vec{V} = ax\hat{i} + bx^2\hat{j} + ax\hat{k}$ (5)  $\vec{V} = ax\hat{i} + [e^{bx}]\hat{j}$  (6)  $\vec{V} = ax\hat{i} + bx^2\hat{j} + ay\hat{k}$ (7)  $\vec{V} = ax\hat{i} + [e^{bt}]\hat{j} + ay\hat{k}$  (8)  $\vec{V} = ax\hat{i} + [e^{by}]\hat{j} + az\hat{k}$ 

# **Given:** Velocity fields

**Find:** Whether flows are 1, 2 or 3D, steady or unsteady.

# Solution:

(1)	$\overrightarrow{V} = \overrightarrow{V}(x,y)$	2D	$\overrightarrow{V} = \overrightarrow{V}(t)$	Unsteady
(2)	$\overrightarrow{V} = \overrightarrow{V}(x,y)$	2D		Steady
(3)	$\overrightarrow{V} = \overrightarrow{V}(x)$	1D	$ \overrightarrow{V} \neq \overrightarrow{V}(t) $	Steady
(4)	$\overrightarrow{V} = \overrightarrow{V}(x)$	1D	$ \overrightarrow{V} \neq \overrightarrow{V}(t) $	Steady
(5)	$\overrightarrow{V} = \overrightarrow{V}(x)$	1D	$\overrightarrow{V} = \overrightarrow{V}(t)$	Unsteady
(6)	$\overrightarrow{V} = \overrightarrow{V}(x,y)$	2D	$ \overrightarrow{V} \neq \overrightarrow{V}(t) $	Steady
(7)	$\overrightarrow{V} = \overrightarrow{V}(x,y)$	2D	$\overrightarrow{V} = \overrightarrow{V}(t)$	Unsteady
(8)	$\overrightarrow{V} = \overrightarrow{V}(x,y,z)$	3D	$\overrightarrow{V} \neq \overrightarrow{V}(t)$	Steady

- a. whether the flow field is one-, two-, or three-dimensional, and why.
- b. whether the flow is steady or unsteady, and why. (The quantities *a* and *b* are constants.)

(1)  $\vec{V}^{\,a} = [ay^2 e^{-bt}]\hat{i}$  (2)  $\vec{V} = ax^2\hat{i} + bx\hat{j} + c\hat{k}$ (3)  $\vec{V} = axy\hat{i} - byt\hat{j}$  (4)  $\vec{V} = ax\hat{i} - by\hat{j} + ct\hat{k}$ (5)  $\vec{V} = [ae^{-bx}]\hat{i} + bt^2\hat{j}$  (6)  $\vec{V} = a(x^2 + y^2)^{1/2}(1/z^3)\hat{k}$ (7)  $\vec{V} = (ax + t)\hat{i} - by^2\hat{j}$  (8)  $\vec{V} = ax^2\hat{i} + bxz\hat{j} + cy\hat{k}$ 

# **Given:** Velocity fields

**Find:** Whether flows are 1, 2 or 3D, steady or unsteady.

# Solution:

(1)	$\overrightarrow{V} = \overrightarrow{V}(y)$	1D	$\overrightarrow{V} = \overrightarrow{V}(t)$	Unsteady
(2)	$\overrightarrow{V} = \overrightarrow{V}(x)$	1D		Steady
(3)	$\overrightarrow{V} = \overrightarrow{V}(x,y)$	2D	$\overrightarrow{V} = \overrightarrow{V}(t)$	Unsteady
(4)	$\overrightarrow{V} = \overrightarrow{V}(x,y)$	2D	$\overrightarrow{V} = \overrightarrow{V}(t)$	Unsteady
(5)	$\overrightarrow{V} = \overrightarrow{V}(x)$	1D	$\overrightarrow{V} = \overrightarrow{V}(t)$	Unsteady
(6)	$\overrightarrow{V} = \overrightarrow{V}(x,y,z)$	3D	$\overrightarrow{V} \neq \overrightarrow{V}(t)$	Steady
(7)	$\overrightarrow{V} = \overrightarrow{V}(x,y)$	2D	$\overrightarrow{V} = \overrightarrow{V}(t)$	Unsteady
(8)	$\overrightarrow{V} = \overrightarrow{V}(x,y,z)$	3D	$\overrightarrow{V} \neq \overrightarrow{V}(t)$	Steady

**2.3** A viscous liquid is sheared between two parallel disks; the upper disk rotates and the lower one is fixed. The velocity field between the disks is given by  $\vec{V} = \hat{e}_0 r \omega z/h$ . (The origin of coordinates is located at the center of the lower disk; the upper disk is located at z = h.) What are the dimensions of this velocity field? Does this velocity field satisfy appropriate physical boundary conditions? What are they?

Given:

Viscous liquid sheared between parallel disks.

Upper disk rotates, lower fixed.

Velocity field is:

# Find:

- a. Dimensions of velocity field.
- b. Satisfy physical boundary conditions.

**Solution:** To find dimensions, compare to  $\vec{V} = \vec{V}(x, y, z)$  form.

The given field is  $\vec{V} = \vec{V}(r, z)$ . Two space coordinates are included, so the field is 2-D.

 $\vec{V} = \hat{e}_{\theta} \frac{r\omega z}{h}$ 

Flow must satisfy the no-slip condition:

1. At lower disk,  $\vec{V} = 0$  since stationary.

$$z = 0$$
, so  $\vec{V} = \hat{e}_{\theta} \frac{r\omega 0}{h} = 0$ , so satisfied.

2. At upper disk,  $\vec{V} = \hat{e}_{\theta} r \omega$  since it rotates as a solid body.

$$z = h$$
, so  $\vec{V} = \hat{e}_{\theta} \frac{r\omega h}{h} = \hat{e}_{\theta} r\omega$ , so satisfied.



**2.4** For the velocity field  $\vec{V} = Ax^2y\hat{i} + Bxy^2\hat{j}$ , where  $A = 2 \text{ m}^{-2}\text{s}^{-1}$  and  $B = 1 \text{ m}^{-2}\text{s}^{-1}$ , and the coordinates are measured in meters, obtain an equation for the flow streamlines. Plot several streamlines in the first quadrant.

Given:	Velocity field	
Find:	Equation for streamlines	
Solution:		Streamline Plots
For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{dy}}{\mathbf{dx}} = \frac{\mathbf{B} \cdot \mathbf{x} \cdot \mathbf{y}^2}{\mathbf{A} \cdot \mathbf{x}^2 \cdot \mathbf{y}} = \frac{\mathbf{B} \cdot \mathbf{y}}{\mathbf{A} \cdot \mathbf{x}}$	$\begin{array}{c} C = 1 \\ C = 2 \\ C = 3 \\ C = 4 \end{array}$
So, separating variables	$\frac{\mathrm{d}y}{\mathrm{y}} = \frac{\mathrm{B}}{\mathrm{A}} \cdot \frac{\mathrm{d}x}{\mathrm{x}}$	$(\underline{\mathbf{u}}) \xrightarrow{3}{7}$
Integrating	$\ln(y) = \frac{B}{A} \cdot \ln(x) + c = -\frac{1}{2} \cdot \ln(x) + c$	
The solution is	$y = \frac{C}{\sqrt{x}}$	
		x (m)

The plot can be easily done in *Excel*.

2.5 The velocity field v interpreted to represe for the flow streamlin several streamlines in that passes through th	$\vec{\nabla} = Ax\hat{i} - Ay\hat{j}$ , where ent flow in a corner. nes. Explain the relet in the first quadrant, is the point $(x, y) = (0, 0)$	$A = 2 \text{ s}^{-1}$ , can be Find an equation evance of A. Plot including the one ).			
Given:	Velocity field				
Find:	Equation for stream	lines; Plot several in the first qu	adrant, including	g one that passes th	hrough point (0,0)
Solution:					
Governing equation:	For streamlines	$\frac{v}{u} = \frac{dy}{dx}$			
Assumption: 2D flow					
Hence		$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}} = -\frac{\mathbf{A}\cdot\mathbf{y}}{\mathbf{A}\cdot\mathbf{x}} = -\frac{\mathbf{y}}{\mathbf{x}}$	51	Streamli	ne Plots
So, separating variables	5	$\frac{\mathrm{d}y}{\mathrm{y}} = -\frac{\mathrm{d}x}{\mathrm{x}}$	4-		C = 1 C = 2 C = 3
Integrating		$\ln(y) = -\ln(x) + c$	<del>ا</del> (آ		C = 4
The solution is		$\ln(\mathbf{x} \cdot \mathbf{y}) = \mathbf{c}$	≻ <sub>2</sub> -		
or		$y = \frac{C}{x}$	r+		
The plot can be easily d	lone in <i>Excel</i> .		0	1 2 x (1	3 4 5 m)

The streamline passing through (0,0) is given by the vertical axis, then the horizontal axis. The value of A is irrelevant to streamline shapes but IS relevant for computing the velocity at each point. **2.6** A velocity field is specified as  $\vec{V} = axy\hat{i} + by^2\hat{j}$ , where  $a = 2 \text{ m}^{-1}\text{s}^{-1}$ ,  $b = -6 \text{ m}^{-1}\text{s}^{-1}$ , and the coordinates are measured in meters. Is the flow field one-, two-, or three-dimensional? Why? Calculate the velocity components at the point (2, ½). Develop an equation for the streamline passing through this point. Plot several streamlines in the first quadrant including the one that passes through the point (2, ½).

**Given:** Velocity field

Find:

Whether field is 1D, 2D or 3D; Velocity components at (2,1/2); Equation for streamlines; Plot

# Solution:

The velocity field is a function of x and y. It is therefore 2D.



This can be plotted in Excel.

**2.7** A velocity field is given by  $\vec{V} = ax\hat{i} - bty\hat{j}$ , where  $a = 1 \text{ s}^{-1}$  and  $b = 1 \text{ s}^{-2}$ . Find the equation of the streamlines at any time *t*. Plot several streamlines in the first quadrant at t = 0 s, t = 1 s, and t = 20 s.

# Given: Velocity field

FING. Equation for streamlines: Plot streamlin
--

# Solution:

For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{d\mathbf{y}}{d\mathbf{x}} = \frac{-\mathbf{b}\cdot\mathbf{t}\cdot\mathbf{y}}{\mathbf{a}\cdot\mathbf{x}}$
So, separating variables	$\frac{\mathrm{d}y}{\mathrm{y}} = \frac{-\mathrm{b}\cdot\mathrm{t}}{\mathrm{a}}\cdot\frac{\mathrm{d}x}{\mathrm{x}}$
Integrating	$\ln(y) = \frac{-b \cdot t}{a} \cdot \ln(x)$
The solution is	$y = c \cdot x^{a}$
For $t = 0$ s $y = c$	For $t = 1$ s $y = \frac{c}{x}$ For $t = 20$ s $y = c \cdot x^{-20}$

### t = 0

	c = 1	<b>c</b> = 2	c = 3
Х	У	У	У
0.05	1.00	2.00	3.00
0.10	1.00	2.00	3.00
0.20	1.00	2.00	3.00
0.30	1.00	2.00	3.00
0.40	1.00	2.00	3.00
0.50	1.00	2.00	3.00
0.60	1.00	2.00	3.00
0.70	1.00	2.00	3.00
0.80	1.00	2.00	3.00
0.90	1.00	2.00	3.00
1.00	1.00	2.00	3.00
1.10	1.00	2.00	3.00
1.20	1.00	2.00	3.00
1.30	1.00	2.00	3.00
1.40	1.00	2.00	3.00
1.50	1.00	2.00	3.00
1.60	1.00	2.00	3.00
1.70	1.00	2.00	3.00
1.80	1.00	2.00	3.00
1.90	1.00	2.00	3.00
2.00	1.00	2.00	3.00

t =1 s				
(### means	too	large	to	view)

	c = 1	<b>c</b> = 2	c = 3
X	У	У	У
0.05	20.00	40.00	60.00
0.10	10.00	20.00	30.00
0.20	5.00	10.00	15.00
0.30	3.33	6.67	10.00
0.40	2.50	5.00	7.50
0.50	2.00	4.00	6.00
0.60	1.67	3.33	5.00
0.70	1.43	2.86	4.29
0.80	1.25	2.50	3.75
0.90	1.11	2.22	3.33
1.00	1.00	2.00	3.00
1.10	0.91	1.82	2.73
1.20	0.83	1.67	2.50
1.30	0.77	1.54	2.31
1.40	0.71	1.43	2.14
1.50	0.67	1.33	2.00
1.60	0.63	1.25	1.88
1.70	0.59	1.18	1.76
1.80	0.56	1.11	1.67
1.90	0.53	1.05	1.58
2.00	0.50	1.00	1.50

t =	20	S
-----	----	---

	c = 1	c = 2	c = 3
Х	У	У	у
0.05	######	######	######
0.10	######	######	######
0.20	######	######	######
0.30	######	######	######
0.40	######	######	######
0.50	######	######	######
0.60	######	######	######
0.70	######	######	######
0.80	86.74	173.47	260.21
0.90	8.23	16.45	24.68
1.00	1.00	2.00	3.00
1.10	0.15	0.30	0.45
1.20	0.03	0.05	0.08
1.30	0.01	0.01	0.02
1.40	0.00	0.00	0.00
1.50	0.00	0.00	0.00
1.60	0.00	0.00	0.00
1.70	0.00	0.00	0.00
1.80	0.00	0.00	0.00
1.90	0.00	0.00	0.00
2.00	0.00	0.00	0.00



**2.8** A velocity field is given by  $\vec{V} = ax^3\hat{i} + bxy^3\hat{j}$ , where  $a = 1 \text{ m}^{-2}\text{s}^{-1}$  and  $b = 1 \text{ m}^{-3}\text{s}^{-1}$ . Find the equation of the streamlines. Plot several streamlines in the first quadrant.

# Given: Velocity field

Find:	Equation for streamlines; Plot str	eamlines
-------	------------------------------------	----------

# Solution:

Streamlines are given by	$\frac{v}{u} = \frac{dy}{dx} = \frac{b \cdot x \cdot y^3}{a \cdot x^3}$
So, separating variables	$\frac{\mathrm{d}y}{\mathrm{y}^3} = \frac{\mathrm{b} \cdot \mathrm{d}x}{\mathrm{a} \cdot \mathrm{x}^2}$
Integrating	$-\frac{1}{2 \cdot y^2} = \frac{b}{a} \cdot \left(-\frac{1}{x}\right) + C$
The solution is	$y = \frac{1}{\sqrt{2 \cdot \left(\frac{b}{a \cdot x} + C\right)}}$

Note: For convenience the sign of C is changed.

a	=	I

n	_	_
	_	

<b>C</b> =	0	2	4	6
Х	У	У	У	У
0.05	0.16	0.15	0.14	0.14
0.10	0.22	0.20	0.19	0.18
0.20	0.32	0.27	0.24	0.21
0.30	0.39	0.31	0.26	0.23
0.40	0.45	0.33	0.28	0.24
0.50	0.50	0.35	0.29	0.25
0.60	0.55	0.37	0.30	0.26
0.70	0.59	0.38	0.30	0.26
0.80	0.63	0.39	0.31	0.26
0.90	0.67	0.40	0.31	0.27
1.00	0.71	0.41	0.32	0.27
1.10	0.74	0.41	0.32	0.27
1.20	0.77	0.42	0.32	0.27
1.30	0.81	0.42	0.32	0.27
1.40	0.84	0.43	0.33	0.27
1.50	0.87	0.43	0.33	0.27
1.60	0.89	0.44	0.33	0.27
1.70	0.92	0.44	0.33	0.28
1.80	0.95	0.44	0.33	0.28
1.90	0.97	0.44	0.33	0.28
2.00	1.00	0.45	0.33	0.28



**2.9** A flow is described by the velocity field  $\vec{V} = (Ax + B)\hat{i} + \hat{i}$  $(-Ay)\hat{j}$ , where A = 10 ft/s/ft and B = 20 ft/s. Plot a few streamlines in the xy plane, including the one that passes through the point (x, y) = (1, 2).

Given: Velocity field

Find: Plot streamlines

### Solution:

Streamlines are given by

So, separating variables

Integrating

The solution is

 $\frac{\mathrm{d} y}{-\mathrm{A} \cdot y} = \frac{\mathrm{d} x}{\mathrm{A} \cdot x + \mathrm{B}}$  $-\frac{1}{A}\ln(y) = \frac{1}{A} \cdot \ln\left(x + \frac{B}{A}\right)$  $y = \frac{C}{x + \frac{B}{A}}$ 

 $\frac{v}{u} = \frac{dy}{dx} = \frac{-A \cdot y}{A \cdot x + B}$ 

$$y = \frac{6}{x+2}$$



$$= \frac{6}{x + \frac{20}{10}}$$

$$\begin{array}{rrr} \mathbf{A}=&\mathbf{10}\\ \mathbf{B}=&\mathbf{20} \end{array}$$

**C** =

	L	4	4	0
X	У	У	У	У
0.00	0.50	1.00	2.00	3.00
0.10	0.48	0.95	1.90	2.86
0.20	0.45	0.91	1.82	2.73
0.30	0.43	0.87	1.74	2.61
0.40	0.42	0.83	1.67	2.50
0.50	0.40	0.80	1.60	2.40
0.60	0.38	0.77	1.54	2.31
0.70	0.37	0.74	1.48	2.22
0.80	0.36	0.71	1.43	2.14
0.90	0.34	0.69	1.38	2.07
1.00	0.33	0.67	1.33	2.00
1.10	0.32	0.65	1.29	1.94
1.20	0.31	0.63	1.25	1.88
1.30	0.30	0.61	1.21	1.82
1.40	0.29	0.59	1.18	1.76
1.50	0.29	0.57	1.14	1.71
1.60	0.28	0.56	1.11	1.67
1.70	0.27	0.54	1.08	1.62
1.80	0.26	0.53	1.05	1.58
1.90	0.26	0.51	1.03	1.54
2.00	0.25	0.50	1.00	1.50



Given:	Velocity field						
Find:	Equation for streamline	ne through (1,3)					
Solution:		$A \cdot \frac{y}{y}$					
For streamlines		$\frac{v}{u} = \frac{dy}{dx} = \frac{x^2}{\underline{A}} =$	$=\frac{y}{x}$				
So, separating variables		$\frac{dy}{y} = \frac{dx}{x}$					
Integrating		$\ln(y) = \ln(x) + c$					
The solution is		$y = C \cdot x$	which is t	he equation of a stra	ight line.		
For the streamline throug	h point (1,3)	$3 = C \cdot 1$	C = 3	and	$y = 3 \cdot x$		
For a particle		$u_p = \frac{dx}{dt} = \frac{A}{x}$	or	$\mathbf{x} \cdot \mathbf{dx} = \mathbf{A} \cdot \mathbf{dt}$	$\mathbf{x} = \sqrt{2 \cdot \mathbf{A} \cdot \mathbf{t} + \mathbf{c}}$	$t = \frac{x^2}{2 \cdot A} -$	$\frac{c}{2 \cdot A}$

Hence the time for a particle to go from x = 1 to x = 2 m is

$$\Delta t = t(x = 2) - t(x = 1) \qquad \Delta t = \frac{(2 \cdot m)^2 - c}{2 \cdot A} - \frac{(1 \cdot m)^2 - c}{2 \cdot A} = \frac{4 \cdot m^2 - 1 \cdot m^2}{2 \times 2 \cdot \frac{m^2}{s}} \qquad \Delta t = 0.75 \cdot s$$

2.11 The flow field for an atmospheric flow is given by

$$\vec{V} = -\frac{My}{2\pi}\hat{i} + \frac{Mx}{2\pi}\hat{j}$$

where  $M = 1 \text{ s}^{-1}$ , and the *x* and *y* coordinates are the parallel to the local latitude and longitude. Plot the velocity magnitude along the *x* axis, along the *y* axis, and along the line y = x, and discuss the velocity direction with respect to these three axes. For each plot use a range *x* or y = 0 km to 1 km. Find the equation for the streamlines and sketch several of them. What does this flow field model?

**Given:** Flow field

**Find:** Plot of velocity magnitude along axes, and y = x; Equation for streamlines

# Solution:



The velocity is perpendicular to the axis and increases linearly with distance x.





y (km)

The velocity is perpendicular to the axis and increases linearly with distance y. This can also be plotted in Excel.

On the $y = y$	M·y M·x	M·x
On the $y - x$	u = =	$V \equiv$
axis	$2\cdot\pi$ $2\cdot\pi$	$2 \cdot \pi$

The flow is perpendicular to line y = x:

x:  
Slope of trajectory of 
$$\frac{u}{v} = -1$$
  
motion:  
If we define the radial position:  
 $r = \sqrt{x^2 + y^2}$  then along  $y = r = \sqrt{x^2 + x^2} = \sqrt{2} \cdot x$   
Then the magnitude of the velocity along  $y = x$  is  $V = \sqrt{u^2 + v^2} = \frac{M}{2 \cdot \pi} \cdot \sqrt{x^2 + x^2} = \frac{M \cdot \sqrt{2} \cdot x}{2 \cdot \pi} = \frac{M \cdot r}{2 \cdot \pi}$ 

 $M\!\cdot\!x$ 

Slope of line y =

Plotting





1

This can also be plotted in Excel.

For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{dy}}{\mathbf{dx}} = \frac{2 \cdot \pi}{-\frac{\mathbf{M} \cdot \mathbf{y}}{2 \cdot \pi}} = -\frac{\mathbf{x}}{\mathbf{y}}$
So, separating variables	$\mathbf{y} \cdot \mathbf{d}\mathbf{y} = -\mathbf{x} \cdot \mathbf{d}\mathbf{x}$
Integrati ng	$\frac{y^2}{2} = -\frac{x^2}{2} + c$
The solution	$x^{2} + y^{2} = C$ which is the equation of a circle.

is The streamlines form a set of concentric circles.

This flow models a rigid body vortex flow. See Example 5.6 for streamline plots. Streamlines are circular, and the velocity approaches zero as we approach the center. In Problem 2.10, we see that the streamlines are also circular. In a real tornado, at large distances from the center, the velocities behave as in Problem 2.10; close to the center, they behave as in this problem.

2.12 The flow field for an atmospheric flow is given by

$$\vec{V} = -\frac{Ky}{2\pi(x^2+y^2)}\hat{i} + \frac{Kx}{2\pi(x^2+y^2)}\hat{j}$$

where  $K = 10^5 \text{ m}^2/\text{s}$ , and the x and y coordinates are parallel to the local latitude and longitude. Plot the velocity magnitude along the x axis, along the y axis, and along the line y = x, and discuss the velocity direction with respect to these three axes. For each plot use a range x or y = -1 km to 1 km, excluding |x|or |y| < 100 m. Find the equation for the streamlines and sketch several of them. What does this flow field model?

Given: Flow field

Find: Plot of velocity magnitude along axes, and y = x; Equation of streamlines

# Solution:



x (km)

The velocity is perpendicular to the axis, is very high close to the origin, and falls off to zero.

This can also be plotted in Excel.

Plotting

 $u = -\frac{K \cdot y}{2 \cdot \pi \cdot \left(x^2 + y^2\right)} = -\frac{K}{2 \cdot \pi \cdot y} \qquad v = \frac{K \cdot x}{2 \cdot \pi \cdot \left(x^2 + y^2\right)} = 0$ On the y axis, x = 0, so 16080 v( m/s) - 0.5 05 0 -80- 160



The velocity is perpendicular to the axis, is very high close to the origin, and falls off to zero. This can also be plotted in Excel.

On the y = x axis  
$$u = -\frac{K \cdot x}{2 \cdot \pi \cdot \left(x^2 + x^2\right)} = -\frac{K}{4 \cdot \pi \cdot x} \qquad v = \frac{K \cdot x}{2 \cdot \pi \cdot \left(x^2 + x^2\right)} = \frac{K}{4 \cdot \pi \cdot x}$$

The flow is perpendicular to line y = x:  
Slope of line y = x:  
Slope of line y = x:  
Slope of trajectory of motion:  

$$\frac{u}{v} = -1$$
  
If we define the radial position:  
 $r = \sqrt{x^2 + y^2}$  then along y = x  
Then the magnitude of the velocity along y = x is  
 $V = \sqrt{u^2 + v^2} = \frac{K}{4 \cdot \pi} \cdot \sqrt{\frac{1}{x^2} + \frac{1}{x^2}} = \frac{K}{2 \cdot \pi \cdot \sqrt{2} \cdot x} = \frac{K}{2 \cdot \pi \cdot \sqrt{2} \cdot x}$   
Plotting  
 $\int_{U}^{U}$   
 $\int_$ 



This can also be plotted in Excel.



So, separating variables

 $y \cdot dy = -x \cdot dx$ 

Integrating

For streamlines

The

 $\frac{y^2}{2} = -\frac{x^2}{2} + c$ 

The solution is

 $x^2 + y^2 = C$ which is the equation of a circle.

Streamlines form a set of concentric circles.

This flow models a vortex flow. See Example 5.6 for streamline plots. Streamlines are circular, and the velocity approaches infinity as we approach the center. In Problem 2.11, we see that the streamlines are also circular. In a real tornado, at large distances from the center, the velocities behave as in this problem; close to the center, they behave as in Problem 2.11.

$$\vec{V} = -\frac{qx}{2\pi(x^2+y^2)}\hat{i} - \frac{qy}{2\pi(x^2+y^2)}\hat{j}$$

where  $q = 5 \times 10^4$  m<sup>2</sup>/s. Plot the velocity magnitude along the x axis, along the y axis, and along the line y = x, and discuss the velocity direction with respect to these three axes. For each plot use a range x or y = -1 km to 1 km, excluding |x| or |y| < 100 m. Find the equation for the streamlines and sketch several of them. What does this flow field model?

### **Given:** Flow field

**Find:** Plot of velocity magnitude along axes, and y = x; Equations of streamlines

# Solution:

Plotting



x (km)

The velocity is very high close to the origin, and falls off to zero. It is also along the axis. This can be plotted in *Excel*.

On the y axis, 
$$x = 0$$
, so  $u = -\frac{q \cdot x}{2 \cdot \pi \cdot \left(x^2 + y^2\right)} = 0$   $v = -\frac{q \cdot y}{2 \cdot \pi \cdot \left(x^2 + y^2\right)} = -\frac{q}{2 \cdot \pi \cdot y}$ 



y (km)

The velocity is again very high close to the origin, and falls off to zero. It is also along the axis.

This can also be plotted in *Excel*.



 $2 \cdot \pi \cdot (x^2 +$ 

The solution is  $y = C \cdot x$  which is the equation of a straight line.

 $\frac{dy}{y} = \frac{dx}{x}$ 

 $\ln(y) = \ln(x) + c$ 

This flow field corresponds to a sink (discussed in Chapter 6).

So, separating variables

Integrating

**2.14** Beginning with the velocity field of Problem 2.5, show that the parametric equations for particle motion are given by  $x_p = c_1 e^{At}$  and  $y_p = c_2 e^{-At}$ . Obtain the equation for the pathline of the particle located at the point (x, y) = (2, 2) at the instant t = 0. Compare this pathline with the streamline through the same point.

# **Given:** Velocity field

**Find:** Proof that the parametric equations for particle motion are  $x_p = c_1 \cdot e^{A \cdot t}$  and  $y_p = c_2 \cdot e^{-A \cdot t}$ ; pathline that was at (2,2) at t = 0; compare to streamline through same point, and explain why they are similar or not.

## Solution:

Governing equations:	For pathlines	$u_p = \frac{dx}{dt}$	$v_p = \frac{dy}{dt}$	For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}}$
Assumption: 2D flow					
Hence for pathlines	$u_p = \frac{dx}{dt} = A \cdot x$			$\mathbf{v}_{\mathbf{p}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{t}} = -\mathbf{A} \cdot \mathbf{y}$	
So, separating variables	$\frac{\mathrm{d}x}{x} = A \cdot \mathrm{d}t$			$\frac{\mathrm{d}y}{\mathrm{y}} = -\mathbf{A} \cdot \mathrm{d}t$	
Integrating	$\ln(\mathbf{x}) = \mathbf{A} \cdot \mathbf{t} + \mathbf{C}_1$			$\ln(y) = -A \cdot t + C_2$	
	$x = e^{A \cdot t + C_1} = e^{C_1}$	$c^{1} \cdot e^{\mathbf{A} \cdot \mathbf{t}} = c_{1} \cdot e^{\mathbf{A} \cdot \mathbf{t}}$	t	$y = e^{-A \cdot t + C_2} = e^{C_2}$	$e^{2} \cdot e^{-A \cdot t} = c_{2} \cdot e^{-A \cdot t}$
The pathlines are	$x = c_1 \cdot e^{A \cdot t}$			$y = c_2 \cdot e^{-A \cdot t}$	
Eliminating t	$t = \frac{1}{A} \cdot \ln\left(\frac{x}{c_1}\right) =$	$-\frac{1}{A} \cdot \ln\left(\frac{y}{c_2}\right)$		$\ln\left(x\frac{1}{x},\frac{1}{y},\frac{1}{x}\right) = \cos x$	t or $\ln(x^A \cdot y^A) = \text{const}$
			SO	$x^{A} \cdot y^{A} = const$ or	$\mathbf{x} \cdot \mathbf{y} = 4$ for given data
For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathrm{d}\mathbf{y}}{\mathrm{d}\mathbf{x}} = -\frac{\mathbf{A}\cdot\mathbf{y}}{\mathbf{A}\cdot\mathbf{x}}$	$=\frac{y}{x}$			
So, separating variables	$\frac{\mathrm{d}y}{\mathrm{y}} = -\frac{\mathrm{d}x}{\mathrm{x}}$				
Integrating	$\ln(y) = -\ln(x) + $	c			
The solution is	$\ln(x \cdot y) = c$	or $x \cdot y =$	= const or	$x \cdot y = 4$ for	or given data

The streamline passing through (2,2) and the pathline that started at (2,2) coincide because the flow is steady!

**2.15** A flow field is given by  $\vec{V} = Ax\hat{i} + 2Ay\hat{j}$ , where  $A = 2 \text{ s}^{-1}$ . Verify that the parametric equations for particle motion are given by  $x_p = c_1 e^{At}$  and  $y_p = c_2 e^{2At}$ . Obtain the equation for the pathline of the particle located at the point (x, y) = (2, 2) at the instant t = 0. Compare this pathline with the streamline through the same point.

### **Given:** Velocity field

**Find:** Proof that the parametric equations for particle motion are  $x_p = c_1 \cdot e^{A \cdot t}$  and  $y_p = c_2 \cdot e^{2 \cdot A \cdot t}$ ; pathline that was at (2,2) at t = 0; compare to streamline through same point, and explain why they are similar or not.

### Solution:

Governing equations:	For pathlines $u_p = \frac{dx}{dt}$	$v_p = \frac{dy}{dt}$	For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}}$
Assumption: 2D flow				
Hence for pathlines	$u_p = \frac{dx}{dt} = A \cdot x$		$v_p = \frac{dy}{dt} = 2 \cdot A$	А·у
So, separating variables	$\frac{\mathrm{d}x}{\mathrm{x}} = \mathrm{A} \cdot \mathrm{d}t$		$\frac{\mathrm{d}y}{\mathrm{y}} = 2 \cdot \mathrm{A} \cdot \mathrm{d}t$	
Integrating	$\ln(\mathbf{x}) = \mathbf{A} \cdot \mathbf{t} + \mathbf{C}_1$		$\ln(\mathbf{y}) = 2 \cdot \mathbf{A} \cdot \mathbf{t} +$	+ C <sub>2</sub>
	$\mathbf{x} = \mathbf{e}^{\mathbf{A} \cdot \mathbf{t} + \mathbf{C}_1} = \mathbf{e}^{\mathbf{C}_1} \cdot \mathbf{e}^{\mathbf{A}_1}$	$^{A \cdot t} = c_1 \cdot e^{A \cdot t}$	$y = e^{2 \cdot A \cdot t + C_2}$	$= e^{C_2} \cdot e^{2 \cdot A \cdot t} = c_2 \cdot e^{2 \cdot A \cdot t}$
The pathlines are	$\mathbf{x} = \mathbf{c}_1 \cdot \mathbf{e}^{\mathbf{A} \cdot \mathbf{t}}$		$y = c_2 \cdot e^{2 \cdot A \cdot t}$	
Eliminating t	$y = c_2 \cdot e^{2 \cdot A \cdot t} = c_2 \cdot \left(\frac{1}{2}\right)$	$\left(\frac{x}{c_1}\right)^2$ so	$y = c \cdot x^2$	or $y = \frac{1}{2} \cdot x^2$ for given data
For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}} = \frac{2 \cdot \mathbf{A} \cdot \mathbf{y}}{\mathbf{A} \cdot \mathbf{x}} = \frac{2}{\mathbf{A} \cdot \mathbf{x}}$	$\frac{2 \cdot y}{x}$		
So, separating variables	$\frac{\mathrm{d}y}{\mathrm{y}} = \frac{2 \cdot \mathrm{d}x}{\mathrm{x}}$	Integrating	ln(y)	$= 2 \cdot \ln(x) + c$
The solution is	$\ln\left(\frac{y}{x^2}\right) = c$			
or	$y = C \cdot x^2$	or	$y = \frac{1}{2} \cdot x^2$ for	or given data

[Difficulty: 2]

The streamline passing through (2,2) and the pathline that started at (2,2) coincide because the flow is steady!

#### Given: Velocity field

Find: Equation of streamlines; Plot streamlines;	streamlines
--	-------------

### Solution:

Streamlines are given by	<u>v</u> =	dy .	-b·x
,	u	đx	a∙y∙t

So, separating variables

 $a \cdot t \cdot y \cdot dy = -b \cdot x \cdot dx$ 

Integrating

 $\frac{1}{2} \cdot \mathbf{a} \cdot \mathbf{t} \cdot \mathbf{y}^2 = -\frac{1}{2} \cdot \mathbf{b} \cdot \mathbf{x}^2 + \mathbf{C}$  $y = \sqrt{C - \frac{b \cdot x^2}{a \cdot t}}$ 

For 
$$t = 0$$
 s  $x = c$ 

The solution is

For 
$$t = 1$$
 s  $y = \sqrt{C - 4 \cdot x^2}$ 

t = 0			
	C = 1	C = 2	C = 3
x	У	У	У
0.00	1.00	2.00	3.00
0.10	1.00	2.00	3.00
0.20	1.00	2.00	3.00
0.30	1.00	2.00	3.00
0.40	1.00	2.00	3.00
0.50	1.00	2.00	3.00
0.60	1.00	2.00	3.00
0.70	1.00	2.00	3.00
0.80	1.00	2.00	3.00
0.90	1.00	2.00	3.00
1.00	1.00	2.00	3.00
1.10	1.00	2.00	3.00
1.20	1.00	2.00	3.00
1.30	1.00	2.00	3.00
1.40	1.00	2.00	3.00
1.50	1.00	2.00	3.00
1.60	1.00	2.00	3.00
1.70	1.00	2.00	3.00
1.80	1.00	2.00	3.00
1.90	1.00	2.00	3.00
2.00	1.00	2.00	3.00

t =1 s			
	C = 1	<b>C</b> = 2	C = 3
x	У	У	У
0.000	1.00	1.41	1.73
0.025	1.00	1.41	1.73
0.050	0.99	1.41	1.73
0.075	0.99	1.41	1.73
0.100	0.98	1.40	1.72
0.125	0.97	1.39	1.71
0.150	0.95	1.38	1.71
0.175	0.94	1.37	1.70
0.200	0.92	1.36	1.69
0.225	0.89	1.34	1.67
0.250	0.87	1.32	1.66
0.275	0.84	1.30	1.64
0.300	0.80	1.28	1.62
0.325	0.76	1.26	1.61
0.350	0.71	1.23	1.58
0.375	0.66	1.20	1.56
0.400	0.60	1.17	1.54
0.425	0.53	1.13	1.51
0.450	0.44	1.09	1.48
0.475	0.31	1.05	1.45
0.500	0.00	1.00	1.41

t = 20 s			
	C = 1	C = 2	C = 3
х	У	У	У
0.00	1.00	1.41	1.73
0.10	1.00	1.41	1.73
0.20	1.00	1.41	1.73
0.30	0.99	1.41	1.73
0.40	0.98	1.40	1.72
0.50	0.97	1.40	1.72
0.60	0.96	1.39	1.71
0.70	0.95	1.38	1.70
0.80	0.93	1.37	1.69
0.90	0.92	1.36	1.68
1.00	0.89	1.34	1.67
1.10	0.87	1.33	1.66
1.20	0.84	1.31	1.65
1.30	0.81	1.29	1.63
1.40	0.78	1.27	1.61
1.50	0.74	1.24	1.60
1.60	0.70	1.22	1.58
1.70	0.65	1.19	1.56
1.80	0.59	1.16	1.53
1.90	0.53	1.13	1.51
2.00	0.45	1.10	1.48

 $y = \sqrt{C - \frac{x^2}{5}}$ 

For t = 20 s



**2.17** Verify that  $x_p = -a\sin(\omega t)$ ,  $y_p = a\cos(\omega t)$  is the equation for the pathlines of particles for the flow field of Problem 2.12. Find the frequency of motion  $\omega$  as a function of the amplitude of motion, a, and K. Verify that  $x_p = -a\sin(\omega t)$ ,  $y_p = a\cos(\omega t)$  is also the equation for the pathlines of particles for the flow field of Problem 2.11, except that  $\omega$  is now a function of M. Plot typical pathlines for both flow fields and discuss the difference.

**Given:** Pathlines of particles

**Find:** Conditions that make them satisfy Problem 2.10 flow field; Also Problem 2.11 flow field; Plot pathlines

### Solution:

ŀ

The given pathlines are  $x_p = -a \cdot \sin(\omega \cdot t)$   $y_p = a \cdot \cos(\omega \cdot t)$ The velocity field of Problem 2.12 is  $u = -\frac{K \cdot y}{2 \cdot \pi \cdot (x^2 + y^2)}$   $v = \frac{K \cdot x}{2 \cdot \pi \cdot (x^2 + y^2)}$ 

If the pathlines are correct we should be able to substitute x<sub>p</sub> and y<sub>p</sub> into the velocity field to find the velocity as a function of time:

$$u = -\frac{K \cdot y}{2 \cdot \pi \cdot \left(x^2 + y^2\right)} = -\frac{K \cdot a \cdot \cos(\omega \cdot t)}{2 \cdot \pi \cdot \left(a^2 \cdot \sin(\omega \cdot t)^2 + a^2 \cdot \cos(\omega \cdot t)^2\right)} = -\frac{K \cdot \cos(\omega \cdot t)}{2 \cdot \pi \cdot a}$$
(1)  
$$v = \frac{K \cdot x}{2 \cdot \pi \cdot \left(x^2 + y^2\right)} = -\frac{K \cdot (-a \cdot \sin(\omega \cdot t))}{2 \cdot \pi \cdot \left(a^2 \cdot \sin(\omega \cdot t)^2 + a^2 \cdot \cos(\omega \cdot t)^2\right)} = -\frac{K \cdot \sin(\omega \cdot t)}{2 \cdot \pi \cdot a}$$
(2)

$$\frac{\mathrm{d}x_p}{\mathrm{d}t} = u \qquad \qquad \frac{\mathrm{d}x_p}{\mathrm{d}t} = v \tag{2.9}$$

$$u = \frac{dx_p}{dt} = -a \cdot \omega \cdot \cos(\omega \cdot t) \qquad \qquad v = \frac{dy_p}{dt} = -a \cdot \omega \cdot \sin(\omega \cdot t) \qquad (3)$$

Comparing Eqs. 1, 2 and 3 
$$u = -a \cdot \omega \cdot \cos(\omega \cdot t) = -\frac{K \cdot \cos(\omega \cdot t)}{2 \cdot \pi \cdot a}$$
  $v = -a \cdot \omega \cdot \sin(\omega \cdot t) = -\frac{K \cdot \sin(\omega \cdot t)}{2 \cdot \pi \cdot a}$ 

Hence we see that 
$$a \cdot \omega = \frac{K}{2 \cdot \pi \cdot a}$$
 or  $\omega = \frac{K}{2 \cdot \pi \cdot a^2}$  for the pathlines to be correct.

Recall that



To plot this in Excel, compute  $x_p$  and  $y_p$  for t ranging from 0 to 60 s, with  $\omega$  given by the above formula. Plot  $y_p$  versus  $x_p$ . Note that outer particles travel much slower!

This is the free vortex flow discussed in Example 5.6

The velocity field of Problem 2.11 is 
$$u = -\frac{M \cdot y}{2 \cdot \pi}$$
  $v = \frac{M \cdot x}{2 \cdot \pi}$ 

If the pathlines are correct we should be able to substitute x<sub>p</sub> and y<sub>p</sub> into the velocity field to find the velocity as a function of time:

$$u = -\frac{M \cdot y}{2 \cdot \pi} = -\frac{M \cdot (a \cdot \cos(\omega \cdot t))}{2 \cdot \pi} = -\frac{M \cdot a \cdot \cos(\omega \cdot t)}{2 \cdot \pi}$$
(4)

$$\mathbf{v} = \frac{\mathbf{M} \cdot \mathbf{x}}{2 \cdot \pi} = \frac{\mathbf{M} \cdot (-\mathbf{a} \cdot \sin(\omega \cdot \mathbf{t}))}{2 \cdot \pi} = -\frac{\mathbf{M} \cdot \mathbf{a} \cdot \sin(\omega \cdot \mathbf{t})}{2 \cdot \pi}$$
(5)

 $u = \frac{dx_p}{dt} = -a \cdot \omega \cdot \cos(\omega \cdot t) \qquad \qquad v = \frac{dy_p}{dt} = -a \cdot \omega \cdot \sin(\omega \cdot t) \qquad (3)$ 

Comparing Eqs. 1, 4 and 5 
$$u = -a \cdot \omega \cdot \cos(\omega \cdot t) = -\frac{M \cdot a \cdot \cos(\omega \cdot t)}{2 \cdot \pi}$$
  $v = -a \cdot \omega \cdot \sin(\omega \cdot t) = -\frac{M \cdot a \cdot \sin(\omega \cdot t)}{2 \cdot \pi}$ 

Hence we see that  $\omega = \frac{M}{2 \cdot \pi}$  for the pathlines to be correct.

The pathlines



To plot this in Excel, compute  $x_p$  and  $y_p$  for t ranging from 0 to 75 s, with  $\omega$  given by the above formula. Plot  $y_p$  versus  $x_p$ . Note that outer particles travel faster!

This is the forced vortex flow discussed in Example 5.6

Note that this is rigid body rotation!

**2.18** Air flows downward toward an infinitely wide horizontal flat plate. The velocity field is given by  $\vec{V} = (ax\hat{i} - ay\hat{j})(2 + \cos \omega t)$ , where  $a = 5 \text{ s}^{-1}$ ,  $\omega = 2\pi \text{ s}^{-1}$ , x and y (measured in meters) are horizontal and vertically upward, respectively, and t is in s. Obtain an algebraic equation for a streamline at t=0. Plot the streamline that passes through point (x, y) = (3, 3) at this instant. Will the streamline change with time? Explain briefly. Show the velocity vector on your plot at the same point and time. Is the velocity vector tangent to the streamline? Explain.

# **Given:** Time-varying velocity field

**Find:** Streamlines at t = 0 s; Streamline through (3,3); velocity vector; will streamlines change with time

# Solution:

For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{d\mathbf{y}}{d\mathbf{x}} = -\frac{\mathbf{a} \cdot \mathbf{y} \cdot (2 + \cos(\omega \cdot \mathbf{t}))}{\mathbf{a} \cdot \mathbf{x} \cdot (2 + \cos(\omega \cdot \mathbf{t}))} = -\frac{\mathbf{y}}{\mathbf{x}}$
At $t = 0$ (actually all times!)	$\frac{\mathrm{d}y}{\mathrm{d}x} = -\frac{y}{x}$
So, separating variables	$\frac{\mathrm{d}y}{\mathrm{y}} = -\frac{\mathrm{d}x}{\mathrm{x}}$
Integrating	$\ln(y) = -\ln(x) + c$
The solution is	$y = \frac{C}{x}$ which is the equation of a hyperbola.
For the streamline through point (3,3)	$C = \frac{3}{3}$ $C = 1$ and $y = \frac{1}{x}$

The streamlines will not change with time since dy/dx does not change with time.



At 
$$t = 0$$
  
 $u = a \cdot x \cdot (2 + \cos(\omega \cdot t)) = 5 \cdot \frac{1}{s} \times 3 \cdot m \times 3$   
 $u = 45 \cdot \frac{m}{s}$   
 $v = -a \cdot y \cdot (2 + \cos(\omega \cdot t)) = 5 \cdot \frac{1}{s} \times 3 \cdot m \times 3$   
 $v = -45 \cdot \frac{m}{s}$ 

The velocity vector is tangent to the curve;

Tangent of curve at (3,3) is 
$$\frac{dy}{dx} = -\frac{y}{x} = -1$$
  
Direction of velocity at (3,3) is  $\frac{v}{u} = -1$ 

This curve can be plotted in Excel.

[Difficulty: 3]

**2.19** Consider the flow described by the velocity field  $\vec{V} = A(1+Bt)\hat{i} + Cty\hat{j}$ , with A = 1 m/s, B = 1 s<sup>-1</sup>, and C = 1 s<sup>-2</sup>. Coordinates are measured in meters. Plot the pathline traced out by the particle that passes through the point (1, 1) at time t = 0. Compare with the streamlines plotted through the same point at the instants t = 0, 1, and 2 s.

# **Given:** Velocity field

**Find:** Plot of pathline traced out by particle that passes through point (1,1) at t = 0; compare to streamlines through same point at the instants t = 0, 1 and 2s

# Solution:

Governing equations:	For pathlines $u_p = \frac{dx}{dt}$	$v_p = \frac{dy}{dt}$ For streamlines $\frac{v}{u} = \frac{dy}{dx}$	
Assumption: 2D flow			
Hence for pathlines	$u_{p} = \frac{dx}{dt} = A \cdot (1 + B \cdot t)$	$A = 1 \cdot \frac{m}{s}$ $B = 1 \cdot \frac{1}{s}$ $v_p = \frac{dy}{dt} = C \cdot t \cdot y$ $C =$	$= 1 \cdot \frac{1}{s^2}$
So, separating variables	$d\mathbf{x} = \mathbf{A} \cdot (1 + \mathbf{B} \cdot \mathbf{t}) \cdot d\mathbf{t}$	$\frac{\mathrm{d}y}{\mathrm{y}} = \mathbf{C} \cdot \mathbf{t} \cdot \mathrm{d}t$	
Integrating	$\mathbf{x} = \mathbf{A} \cdot \left( \mathbf{t} + \mathbf{B} \cdot \frac{\mathbf{t}^2}{2} \right) + \mathbf{C}_1$	$\ln(\mathbf{y}) = \frac{1}{2} \cdot \mathbf{C} \cdot \mathbf{t}^2 + \mathbf{C}_2$	
		$y = e^{\frac{1}{2} \cdot C \cdot t^{2} + C_{2}} = e^{C_{2}} \cdot e^{\frac{1}{2} \cdot C \cdot t^{2}} = c_{2}$	$\frac{1}{2} \cdot \mathbf{C} \cdot \mathbf{t}^2$
The pathlines are	$\mathbf{x} = \mathbf{A} \cdot \left( \mathbf{t} + \mathbf{B} \cdot \frac{\mathbf{t}^2}{2} \right) + \mathbf{C}_1$	$y = c_2 \cdot e^{\frac{1}{2} \cdot C \cdot t^2}$	
Using given data	$\mathbf{x} = \mathbf{A} \cdot \left( \mathbf{t} + \mathbf{B} \cdot \frac{\mathbf{t}^2}{2} \right) + 1$	$y = e^{\frac{1}{2} \cdot C \cdot t^2}$	
For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}} = \frac{\mathbf{C} \cdot \mathbf{y} \cdot \mathbf{t}}{\mathbf{A} \cdot (1 + \mathbf{B} \cdot \mathbf{t})}$		
So, separating variables	$(1 + B \cdot t) \cdot \frac{dy}{y} = \frac{C}{A} \cdot t \cdot dx$	which we can integrate for any given t (t is treated as a co	nstant)
Integrating	$(1 + B \cdot t) \cdot \ln(y) = \frac{C}{A} \cdot t \cdot x + c$		
The solution is	$y^{1+B\cdot t} = \frac{C}{A} \cdot t \cdot x + const$	$y = \left(\frac{C}{A} \cdot t \cdot x + const\right)^{(1+B \cdot t)}$	





Streamline and Pathline Plots

2.20 Consider the flow described by the velocity field

 $\vec{V} = Bx(1+At)\hat{i} + Cy\hat{j}$ , with  $A = 0.5 \text{ s}^{-1}$  and  $B = C = 1 \text{ s}^{-1}$ . Coordinates are measured in meters. Plot the pathline traced out by the particle that passes through the point (1, 1) at time t = 0. Compare with the streamlines plotted through the same point at the instants t = 0, 1, and 2 s.Given: Velocity field Find: Plot of pathline traced out by particle that passes through point (1,1) at t = 0; compare to streamlines through same point at the instants t = 0, 1 and 2s Solution:  $u_p = \frac{dx}{dt}$   $v_p = \frac{dy}{dt}$  $\frac{v}{u} = \frac{dy}{dx}$ For streamlines **Governing equations:** For pathlines Assumption: 2D flow  $u_p = \frac{dx}{dt} = B \cdot x \cdot (1 + A \cdot t) \qquad A = 0.5 \cdot \frac{1}{s} \qquad B = 1 \cdot \frac{1}{s} \qquad v_p = \frac{dy}{dt} = C \cdot y \qquad C = 1 \cdot \frac{1}{s}$ Hence for pathlines  $\frac{\mathrm{dx}}{\mathrm{x}} = \mathrm{B} \cdot (1 + \mathrm{A} \cdot \mathrm{t}) \cdot \mathrm{dt}$  $\frac{dy}{v} = C \cdot dt$ So, separating variables  $\ln(x) = B \cdot \left( t + A \cdot \frac{t^2}{2} \right) + C_1$  $\ln(\mathbf{y}) = \mathbf{C} \cdot \mathbf{t} + \mathbf{C}_2$ Integrating  $\mathbf{x} = \mathbf{e}^{\mathbf{B} \cdot \left(t + \mathbf{A} \cdot \frac{t^2}{2}\right) + C_1} = \mathbf{e}^{\mathbf{C}_1} \cdot \mathbf{e}^{\mathbf{B} \cdot \left(t + \mathbf{A} \cdot \frac{t^2}{2}\right)} = \mathbf{e}^{\mathbf{C} \cdot \mathbf{e}} \mathbf{B} \cdot \left(t + \mathbf{A} \cdot \frac{t^2}{2}\right)$  $y = e^{C \cdot t + C_2} = e^{C_2} \cdot e^{C \cdot t} = c_2 \cdot e^{C \cdot t}$  $B \cdot \left( t + A \cdot \frac{t^2}{2} \right)$  $x = c_1 \cdot e^{-B \cdot \left( t + A \cdot \frac{t^2}{2} \right)}$  $y = c_2 \cdot e^{C \cdot t}$ The pathlines are  $B \cdot \left( t + A \cdot \frac{t^2}{2} \right)$  $v = e^{C \cdot t}$ Using given data  $\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{dy}}{\mathbf{dx}} = \frac{\mathbf{C} \cdot \mathbf{y}}{\mathbf{B} \cdot \mathbf{x} \cdot (1 + \mathbf{A} \cdot \mathbf{t})}$ For streamlines  $(1 + A \cdot t) \cdot \frac{dy}{v} = \frac{C}{B} \cdot \frac{dx}{x}$ which we can integrate for any given t (t is treated as a constant) So, separating variables  $(1 + A \cdot t) \cdot \ln(y) = \frac{C}{R} \cdot \ln(x) + c$ Integrating





**2.21** Consider the flow field given in Eulerian description by the expression  $\vec{V} = A\hat{i} - Bt\hat{j}$ , where A = 2 m/s, B = 2 m/s<sup>2</sup>, and the coordinates are measured in meters. Derive the Lagrangian position functions for the fluid particle that was located at the point (x, y) = (1, 1) at the instant t = 0. Obtain an algebraic expression for the pathline followed by this particle. Plot the pathline and compare with the streamlines plotted through the same point at the instants t = 0, 1,and 2 s.

Given:

Eulerian V	elocity field						
Lagrangian position function that was at point $(1,1)$ at $t = 0$ ; expression for pathline; plot pathline and compare to streamlines through same point at the instants $t = 0$ , 1 and 2s							
ations:	For pathlines (Lagrangian des	cription) $u_p = \frac{dx}{dt}$	$v_p = \frac{dy}{dt}$ For street	eamlines $\frac{v}{u} = \frac{dy}{dx}$			
2D flow							
ines	$u_p = \frac{dx}{dt} = A$	$A = 2  \frac{m}{s}$	$v_p = \frac{dy}{dt} = -B \cdot t$	$B = 2  \frac{m}{s^2}$			
variables	$d\mathbf{x} = \mathbf{A} \cdot d\mathbf{t}$		$dy = -B \cdot t \cdot dt$				
	$\mathbf{x} = \mathbf{A} \cdot \mathbf{t} + \mathbf{x}_0$	$x_0 = 1 m$	$y = -B \cdot \frac{t^2}{2} + y_0$	$y_0 = 1 m$			
description	is	$\mathbf{x}(t) = \mathbf{A} \cdot \mathbf{t} + \mathbf{x}_0$	$\mathbf{y}(t) = -\mathbf{B} \cdot \frac{t^2}{2} + \mathbf{y}_0$				
a		$\mathbf{x}(\mathbf{t}) = 2 \cdot \mathbf{t} + 1$	$y(t) = 1 - t^2$				
re given by c	combining the equations $t = -$	$\frac{x-x_0}{A}$	$\mathbf{y} = -\mathbf{B} \cdot \frac{\mathbf{t}^2}{2} + \mathbf{y}_0 = -\mathbf{E}$	$3 \cdot \frac{\left(x - x_0\right)^2}{2 \cdot A^2} + y_0$			
	$y(x) = y_0 - B \cdot \frac{(x - x_0)^2}{2 \cdot A^2}$	or, using given data	$y(x) = 1 - \frac{(x-1)^2}{4}$	-			
	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}} = \frac{-\mathbf{B}\cdot\mathbf{t}}{\mathbf{A}}$						
	Eulerian V Lagrangiar streamlines ations: D flow ines description a re given by o	Eulerian Velocity field Lagrangian position function that was at p streamlines through same point at the insta- ations: For pathlines (Lagrangian des- 2D flow ines $u_p = \frac{dx}{dt} = A$ variables $dx = A \cdot dt$ $x = A \cdot t + x_0$ description is a pre given by combining the equations $t = \frac{2}{a}$ $y(x) = y_0 - B \cdot \frac{(x - x_0)^2}{2 \cdot A^2}$ $\frac{v}{u} = \frac{dy}{dx} = \frac{-B \cdot t}{A}$	Eulerian Velocity field Lagrangian position function that was at point (1,1) at t = 0; expressis streamlines through same point at the instants t = 0, 1 and 2s ations: For pathlines (Lagrangian description) $u_p = \frac{dx}{dt}$ D flow ines $u_p = \frac{dx}{dt} = A$ $A = 2 \frac{m}{s}$ variables $dx = A \cdot dt$ $x = A \cdot t + x_0$ $x_0 = 1 \text{ m}$ description is $x(t) = A \cdot t + x_0$ a $x(t) = 2 \cdot t + 1$ re given by combining the equations $t = \frac{x - x_0}{A}$ $y(x) = y_0 - B \cdot \frac{(x - x_0)^2}{2 \cdot A^2}$ or, using given data $\frac{v}{u} = \frac{dy}{dx} = \frac{-B \cdot t}{A}$	Eulerian Velocity field Lagrangian position function that was at point (1,1) at t = 0; expression for pathline; plot pat streamlines through same point at the instants t = 0, 1 and 2s ations: For pathlines (Lagrangian description) $u_p = \frac{dx}{dt}$ $v_p = \frac{dy}{dt}$ For streamlines the for pathlines (Lagrangian description) $u_p = \frac{dx}{dt}$ $v_p = \frac{dy}{dt}$ For streamlines the for pathlines (Lagrangian description) $u_p = \frac{dx}{dt}$ $v_p = \frac{dy}{dt}$ For streamlines the for pathlines (Lagrangian description) $u_p = \frac{dx}{dt}$ $v_p = \frac{dy}{dt} = -B \cdot t$ the formula $u_p = \frac{dx}{dt} = A$ $A = 2$ $\frac{m}{s}$ $v_p = \frac{dy}{dt} = -B \cdot t$ ariables $dx = A \cdot dt$ $dy = -B \cdot t \cdot dt$ $x = A \cdot t + x_0$ $x_0 = 1$ m $y = -B \cdot \frac{t^2}{2} + y_0$ description is $x(t) = A \cdot t + x_0$ $y(t) = -B \cdot \frac{t^2}{2} + y_0$ a $x(t) = 2 \cdot t + 1$ $y(t) = 1 - t^2$ re given by combining the equations $t = \frac{x - x_0}{A}$ $y = -B \cdot \frac{t^2}{2} + y_0 = -B$ $y(x) = y_0 - B \cdot \frac{(x - x_0)^2}{2 \cdot A^2}$ or, using given data $y(x) = 1 - \frac{(x - 1)^2}{4}$ $\frac{v}{u} = \frac{dy}{dx} = \frac{-B \cdot t}{A}$			

So, separating variables

 $dy = -\frac{B \cdot t}{A} \cdot dx$ 

which we can integrate for any given t (t is treated as a constant)

The solution is

$$y = -\frac{B \cdot t}{A} \cdot x + c$$
 and for the one through (1,1)  $1 = -\frac{B \cdot t}{A} \cdot 1 + c$   $c = 1 + \frac{B \cdot t}{A}$ 

$$y = -\frac{B \cdot t}{A} \cdot (x - 1) + 1$$
  $y = 1 - t \cdot (x - 1)$   
 $x = 1, 1.1..20$ 



x (m)

**2.22** Consider the velocity field  $V = ax\hat{i} + by(1 + ct)\hat{j}$ , where  $a = b = 2 \text{ s}^{-1}$  and  $c = 0.4 \text{ s}^{-1}$ . Coordinates are measured in meters. For the particle that passes through the point (x, y) = (1, 1) at the instant t = 0, plot the pathline during the interval from t = 0 to 1.5 s. Compare this pathline with the streamlines plotted through the same point at the instants t = 0, 1, and 1.5 s.

**Given:** Velocity field

**Find:** Plot of pathline of particle for t = 0 to 1.5 s that was at point (1,1) at t = 0; compare to streamlines through same point at the instants t = 0, 1 and 1.5 s

# Solution:

Governing equations:	For pathlines	$u_p = \frac{dx}{dt}$	$v_p = \frac{dy}{dt}$	For streamlines	$\frac{v}{u} = \frac{dy}{dx}$
Assumption: 2D flow					
Hence for pathlines	$u_p = \frac{dx}{dt} = ax$	$a = 2 \frac{1}{s}$	$v_p = \frac{dy}{dt}$	$= \mathbf{b} \cdot \mathbf{y} \cdot (1 + \mathbf{c} \cdot \mathbf{t})  \mathbf{b} = 2$	$\frac{1}{s^2}$ c = 0.4 $\frac{1}{s}$
So, separating variables	$\frac{\mathrm{d}x}{\mathrm{x}} = \mathrm{a} \cdot \mathrm{d}t$		$dy = b \cdot y \cdot$	$(1 + c \cdot t) \cdot dt$ $\frac{dy}{y} = b \cdot (t)$	$(1 + c \cdot t) \cdot dt$
Integrating	$\ln\left(\frac{x}{x_0}\right) = a \cdot t$	$x_0 = 1 m$	$\ln\left(\frac{y}{y_0}\right) =$	$b \cdot \left(t + \frac{1}{2} \cdot c \cdot t^2\right)  y_0 =$	1 m
Hence	$\mathbf{x}(t) = \mathbf{x}_0 \cdot \mathbf{e}^{\mathbf{a} \cdot \mathbf{t}}$		$y(t) = e^{b}$	$\left(t+\frac{1}{2}\cdot c\cdot t^2\right)$	
Using given data	$x(t) = e^{2 \cdot t}$		$y(t) = e^{2}$	$2 \cdot t + 0.4 \cdot t^2$	
For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathrm{d}\mathbf{y}}{\mathrm{d}\mathbf{x}} = \frac{\mathbf{b} \cdot \mathbf{y} \cdot (1 + \mathbf{c} \cdot \mathbf{t})}{\mathbf{a} \cdot \mathbf{x}}$				
So, separating variables	$\frac{\mathrm{d}y}{\mathrm{y}} = \frac{\mathrm{b} \cdot (1 + \mathrm{c} \cdot \mathrm{t})}{\mathrm{a} \cdot \mathrm{x}} \cdot \mathrm{d}\mathrm{x}$	which we can	i integrate for a	ny given t (t is treated as a	constant)
Hence	$\ln\left(\frac{y}{y_0}\right) = \frac{b}{a} \cdot (1 + c \cdot t) \cdot \ln\left(\frac{b}{a}\right)$	$\left(\frac{x}{x_0}\right)$			
The solution is	$y = y_0 \cdot \left(\frac{x}{x_0}\right)^a (1 + c \cdot t)$				

For 
$$t = 0$$
  $y = y_0 \cdot \left(\frac{x}{x_0}\right)^a = x$   $t = 1$   $y = y_0 \cdot \left(\frac{x}{x_0}\right)^a = x^{1.4}$   $t = 1.5$   $y = y_0 \cdot \left(\frac{x}{x_0}\right)^a = x^{1.6}$ 



<b>2.23</b> Consid criptionby th $b = 0.04 \text{ s}^{-2}$ , Derive the L ticle that w the instant t pathline folk compare wit point at the i	er the flo e expressio and the co- agrangian as located r = 0. Obtai owed by the h the strea- instants $t =$	by field given in Euler on $\vec{V} = ax\hat{i} + byt\hat{j}$ , where $a =$ coordinates are measured in position functions for the final at the point $(x, y) = ($ in an algebraic expression his particle. Plot the path amlines plotted through t 0, 10, and 20 s.	ian des- = $0.2 \text{ s}^{-1}$ , n meters. fluid par- 1, 1) at n for the line and he same			
Given:	Velocity	field				
Find:	Plot of pa point at th	thline of particle for $t = 0$ to ne instants $t = 0, 1$ and 1.5 s	o 1.5 s that was at po	int (1,1) at t = 0;	compare to streamlines thr	ough same
Solution:						
Governing equ	uations:	For pathlines	$u_p = \frac{dx}{dt}$	$v_p = \frac{dy}{dt}$	For streamlines	$\frac{v}{u} = \frac{dy}{dx}$
Assumption:	2D flow					
Hence for path	lines	$u_p = \frac{dx}{dt} = a \cdot x$	$a = \frac{1}{5} \frac{1}{s}$	$v_p = \frac{dy}{dt} =$	$b \cdot y \cdot t$ $b = \frac{1}{25} \frac{1}{s}$	<u>1</u> 2
So, separating	variables	$\frac{\mathrm{d}x}{\mathrm{x}} = \mathrm{a} \cdot \mathrm{d}t$		$dy = b \cdot y \cdot t \cdot$	dt $\frac{\mathrm{d}y}{\mathrm{y}} = \mathrm{b}\cdot\mathrm{t}\cdot\mathrm{d}$	lt
Integrating		$\ln\left(\frac{x}{x_0}\right) = a \cdot t$	$x_0 = 1 m$	$ln\left(\frac{y}{y_0}\right) =$	$\mathbf{b} \cdot \frac{1}{2} \cdot \mathbf{t}^2 \qquad \mathbf{y}_0 = 1$	m
Hence		$\mathbf{x}(t) = \mathbf{x}_0 \cdot \mathbf{e}^{\mathbf{a} \cdot \mathbf{t}}$		$\mathbf{y}(t) = \mathbf{y}_0.$	$e^{\frac{1}{2}\cdot\mathbf{b}\cdot\mathbf{t}^2}$	
Using given da	ta	$x(t) = e^{\frac{t}{5}}$		$y(t) = e^{\frac{t^2}{50}}$	2 0	
For streamlines	5	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}} = \frac{\mathbf{b} \cdot \mathbf{y} \cdot \mathbf{t}}{\mathbf{a} \cdot \mathbf{x}}$				
So, separating	variables	$\frac{\mathrm{d}y}{\mathrm{y}} = \frac{\mathrm{b} \cdot \mathrm{t}}{\mathrm{a} \cdot \mathrm{x}} \cdot \mathrm{d}\mathrm{x}$	which we can	n integrate for an	y given t (t is treated as a co	onstant)
Hence		$\ln\left(\frac{y}{y_0}\right) = \frac{b}{a} \cdot t \cdot \ln\left(\frac{x}{x_0}\right)$	) /			
The solution is		$y = y_0 \cdot \left(\frac{x}{x_0}\right)^{\frac{b}{a} \cdot t}$	b a	= 0.2	$x_0 = 1$ $y_0 = 1$	

For

t =

0 
$$y = y_0 \cdot \left(\frac{x}{x_0}\right)^a = 1$$

t = 5 
$$y = y_0 \cdot \left(\frac{x}{x_0}\right)^a = x \qquad \frac{b}{a} \cdot t = 1$$
  
t = 10  $y = y_0 \cdot \left(\frac{x}{x_0}\right)^a = x^2 \qquad \frac{b}{a} \cdot t = 2$ 

Streamline and Pathline Plots



# Given: Velocity field

Find: Plot pathlines and streamlines

### Solution:

Pathlines are given by	$\frac{\mathrm{d}x}{\mathrm{d}t} = u = a \cdot x \cdot t$	$\frac{\mathrm{d}y}{\mathrm{d}t} = v = -b \cdot y$	
So, separating variables	$\frac{\mathrm{d}x}{\mathrm{x}} = \mathbf{a} \cdot \mathbf{t} \cdot \mathrm{d}\mathbf{t}$	$\frac{dy}{y} = -b \cdot dt$	
Integrating	$\ln(\mathbf{x}) = \frac{1}{2} \cdot \mathbf{a} \cdot \mathbf{t}^2 + \mathbf{c}_1$	$\ln(y) = -b \cdot t + c_2$	
For initial position (x <sub>0</sub> ,y <sub>0</sub> )	$x = x_0 \cdot e^{\frac{a}{2} \cdot t^2}$	$y = y_0 \cdot e^{-b \cdot t}$	
Using the given data, and IC $(x_0, y_0) = ($	(1,1) at t = 0		
	$x = e^{0.05 \cdot t^2}$	$y = e^{-t}$	
Streamlines are given by	$\frac{v}{u} = \frac{dy}{dx} = \frac{-b \cdot y}{a \cdot x \cdot t}$		
So, separating variables	$\frac{\mathrm{d}y}{\mathrm{y}} = -\frac{\mathrm{b}}{\mathrm{a}\cdot\mathrm{t}}\cdot\frac{\mathrm{d}x}{\mathrm{x}}$	Integrating	$ln(y) = -\frac{b}{a \cdot t} \cdot ln(x) + C$
The solution is	$y = C \cdot x - \frac{b}{a \cdot t}$		
For streamline at $(1,1)$ at $t = 0$ s	x = c		
For streamline at $(1,1)$ at $t=1$ s	$y = x^{-10}$		

For streamline at (1,1) at t = 2 s  $y = x^{-5}$ 

Pathline **Streamlines** t = 0 t = 2 st = 1 st Х у Х У Х У Х У 0.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 1.00 0.25 1.00 0.78 1.00 0.78 1.00 0.97 1.00 0.98 0.50 1.01 0.61 1.00 0.61 1.01 0.88 1.01 0.94 0.75 1.03 0.47 1.00 0.47 1.03 0.75 1.03 0.87 1.00 1.05 0.37 1.00 0.37 1.05 0.61 1.05 0.78 1.25 1.08 0.29 1.00 0.29 1.08 0.46 1.08 0.68 1.50 1.12 0.22 1.00 0.22 1.12 0.32 1.12 0.57 1.75 1.17 1.00 0.17 0.22 1.17 0.47 0.17 1.17 2.00 1.22 0.14 1.00 0.14 1.22 0.14 1.22 0.37 2.25 1.29 0.11 1.00 0.11 1.29 0.08 1.29 0.28 2.50 1.37 1.00 1.37 0.04 1.37 0.08 0.08 0.21 2.75 1.46 0.06 1.00 0.06 1.46 0.02 1.46 0.15 3.00 1.57 0.05 1.00 0.05 1.57 0.01 1.57 0.11 3.25 1.70 1.70 0.04 1.00 0.04 0.01 1.70 0.07 3.50 1.85 0.03 1.00 0.03 1.85 0.00 1.85 0.05 3.75 2.02 1.00 0.02 2.02 0.00 2.02 0.02 0.03 4.00 2.23 0.02 1.00 0.02 2.23 0.00 2.23 0.02 4.25 2.47 1.00 0.01 2.47 0.00 2.47 0.01 0.01 2.75 2.75 0.00 4.50 0.01 1.00 0.01 2.75 0.01 4.75 3.09 0.01 1.00 0.01 3.09 0.00 3.09 0.00 5.00 3.49 0.01 1.00 0.01 3.49 0.00 3.49 0.00



**2.25** Consider the flow field  $\vec{V} = axt\hat{i} + b\hat{j}$ , where  $a = 0.1 \text{ s}^{-2}$  and b = 4 m/s. Coordinates are measured in meters. For the particle that passes through the point (x, y) = (3, 1) at the instant t = 0, plot the pathline during the interval from t = 0 to 3 s. Compare this pathline with the streamlines plotted through the same point at the instant t = 1, 2, and 3 s.

# **Given:** Flow field

**Find:** Pathline for particle starting at (3,1); Streamlines through same point at t = 1, 2, and 3 s

# Solution:

For particle paths	$\frac{\mathrm{d}x}{\mathrm{d}t} = \mathbf{u} = \mathbf{a} \cdot \mathbf{x} \cdot \mathbf{t}$	an d	$\frac{dy}{dt} = v = b$
Separating variables and integrating	$\frac{\mathrm{d}x}{x} = a \cdot t \cdot \mathrm{d}t$	or	$\ln(x) = \frac{1}{2} \cdot a \cdot t^2 + c_1$
	$dy = b \cdot dt$	or	$y = b \cdot t + c_2$

Using initial condition (x,y) = (3,1) and the given values for a and b

	$c_1 = \ln(3 \cdot m)$	an d	$c_2 = 1 \cdot m$
The pathline is then	$x = 3 \cdot e^{0.05 \cdot t^2}$	and	$y = 4 \cdot t + 1$
For streamlines (at any time t)	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{\mathbf{d}\mathbf{y}}{\mathbf{d}\mathbf{x}} = \frac{\mathbf{b}}{\mathbf{a}\cdot\mathbf{x}\cdot\mathbf{t}}$		
So, separating variables	$dy = \frac{b}{a \cdot t} \cdot \frac{dx}{x}$		
Integrating	$y = \frac{b}{a \cdot t} \cdot \ln(x) + c$		

We are interested in instantaneous streamlines at various times that always pass through point (3,1). Using a and b values:

$$c = y - \frac{b}{a \cdot t} \cdot \ln(x) = 1 - \frac{4}{0.1 \cdot t} \cdot \ln(3)$$
$$y = 1 + \frac{40}{t} \cdot \ln\left(\frac{x}{2}\right)$$

The streamline equation is



These curves can be plotted in Excel.

**2.26** Consider the garden hose of Fig. 2.5. Suppose the velocity field is given by  $\vec{V} = u_0 \hat{i} + v_0 \sin[\omega(t - x/u_0)]\hat{j}$ , where the x direction is horizontal and the origin is at the mean position of the hose,  $u_0 = 10$  m/s,  $v_0 = 2$  m/s, and  $\omega = 5$  cycle/s. Find and plot on one graph the instantaneous streamlines that pass through the origin at t = 0 s, 0.05 s, 0.1 s, and 0.15 s. Also find and plot on one graph the pathlines of particles that left the origin at the same four times.

**Given:** Velocity field

**Find:** Plot streamlines that are at origin at various times and pathlines that left origin at these times

# Solution:

For streamlines	$\frac{v}{u} = \frac{dy}{dx} = \frac{v_0 \cdot \sin\left[\omega \cdot \left(t - \frac{x}{u_0}\right)\right]}{u_0}$	
So, separating variables (t=const)	$dy = \frac{v_0 \cdot \sin \left[\omega \cdot \left(t - \frac{x}{u_0}\right)\right]}{u_0} \cdot dx$	
Integrating	$y = \frac{v_0 \cdot \cos\left[\omega \cdot \left(t - \frac{x}{u_0}\right)\right]}{\omega} + c$	
Using condition $y = 0$ when $x = 0$	$y = \frac{v_0 \cdot \left[ \cos \left[ \omega \cdot \left( t - \frac{x}{u_0} \right) \right] - \cos(\omega \cdot t) \right]}{\omega}$	This gives streamlines $y(x)$ at each time t
For particle paths, first find x(t)	$\frac{dx}{dt} = u = u_0$	
Separating variables and integrating	$dx = u_0 \cdot dt$ o r	$\mathbf{x} = \mathbf{u}_0 \cdot \mathbf{t} + \mathbf{c}_1$
Using initial condition $x = 0$ at $t = \tau$	$c_1 = -u_0 \cdot \tau$	$x = u_0 \cdot (t - \tau)$
For y(t) we have	$\frac{dy}{dt} = v = v_0 \cdot sin\left[\omega \cdot \left(t - \frac{x}{u_0}\right)\right]  so$	$\frac{dy}{dt} = v = v_0 \cdot \sin \left[ \omega \cdot \left[ t - \frac{u_0 \cdot (t - \tau)}{u_0} \right] \right]$
and	$\frac{\mathrm{d}y}{\mathrm{d}t} = v = v_0 \cdot \sin(\omega \cdot \tau)$	
Separating variables and integrating	$dy = v_0 \cdot \sin(\omega \cdot \tau) \cdot dt$	$y = v_0 \cdot \sin(\omega \cdot \tau) \cdot t + c_2$
Using initial condition $y = 0$ at $t = \tau$	$c_2 = -v_0 \cdot \sin(\omega \cdot \tau) \cdot \tau$	$y = v_0 \cdot \sin(\omega \cdot \tau) \cdot (t - \tau)$

The pathline is then

 $x(t,\tau) = u_0 \cdot (t-\tau) \qquad \qquad y(t,\tau) = v_0 \cdot \sin(\omega \cdot \tau) \cdot (t-\tau)$ 

These terms give the path of a particle (x(t),y(t)) that started at  $t = \tau$ .



The streamlines are sinusoids; the pathlines are straight (once a water particle is fired it travels in a straight line). These curves can be plotted in *Excel*.

**2.27** Using the data of Problem 2.26, find and plot the streakline shape produced after the first second of flow.

**Given:** Velocity field

**Find:** Plot streakline for first second of flow

### Solution:

Following the discussion leading up to Eq. 2.10, we first find equations for the pathlines in form

$$x_{p}(t) = x(t, x_{0}, y_{0}, t_{0})$$
 and  $y_{p}(t) = y(t, x_{0}, y_{0}, t_{0})$ 

where  $x_0$ ,  $y_0$  is the position of the particle at  $t = t_0$ , and re-interpret the results as streaklines

$$x_{st}(t_0) = x(t, x_0, y_0, t_0) \qquad \text{and} \qquad y_{st}(t_0) = y(t, x_0, y_0, t_0)$$

which gives the streakline at t, where  $x_0$ ,  $y_0$  is the point at which dye is released ( $t_0$  is varied from 0 to t)

 $\frac{\mathrm{dx}}{\mathrm{dt}} = \mathrm{u} = \mathrm{u}_{0}$ For particle paths, first find x(t)Separating variables and integrating  $dx = u_0 \cdot dt$  $\mathbf{x} = \mathbf{x}_0 + \mathbf{u}_0 \cdot \left( \mathbf{t} - \mathbf{t}_0 \right)$ 0  $\frac{dy}{dt} = v = v_0 \cdot \sin \left[ \omega \cdot \left( t - \frac{x}{u_0} \right) \right] \quad \text{so} \quad \frac{dy}{dt} = v = v_0 \cdot \sin \left[ \omega \cdot \left[ t - \frac{x_0 + u_0 \cdot \left( t - t_0 \right)}{u_0} \right] \right]$ For y(t) we have  $\frac{\mathrm{d}y}{\mathrm{d}t} = v = v_0 \cdot \sin \left[ \omega \cdot \left( t_0 - \frac{x_0}{u_0} \right) \right]$ and Separating variables and integrating  $\mathbf{y}_{st}(t_0) = \mathbf{y}_0 + \mathbf{v}_0 \cdot \sin \left[ \omega \cdot \left( t_0 - \frac{\mathbf{x}_0}{\mathbf{u}_0} \right) \right] \cdot \left( t - t_0 \right)$  $x_{st}(t_0) = x_0 + u_0(t - t_0)$ The streakline is then With  $x_0 = y_0 = 0$  $\mathbf{x}_{st}(t_0) = \mathbf{u}_0 \cdot (t - t_0)$  $y_{st}(t_0) = v_0 \cdot \sin[\omega \cdot (t_0)] \cdot (t - t_0)$ Streakline for First Second y (m) 10

This curve can be plotted in *Excel*. For t = 1,  $t_0$  ranges from 0 to t.

**2.28** Consider the velocity field of Problem 2.20. Plot the streakline formed by particles that passed through the point (1, 1) during the interval from t = 0 to t = 3 s. Compare with the streamlines plotted through the same point at the instants t = 0, 1, and 2 s.

### **Given:** Velocity field

Find: Plot of streakline for t = 0 to 3 s at point (1,1); compare to streamlines through same point at the instants t = 0, 1 and 2 s

# Solution:

Governing equations:	For pathlines	$u_p = \frac{dx}{dt}$	$v_p = \frac{dy}{dt}$	For streamlines	$\frac{\mathbf{v}}{\mathbf{u}} = \frac{1}{2}$	$\frac{dy}{dx}$
		<sup>1</sup> dt	<sup>1</sup> at		u i	ax

Following the discussion leading up to Eq. 2.10, we first find equations for the pathlines in form

$$\begin{aligned} x_{p}(t) &= x \Big( t, x_{0}, y_{0}, t_{0} \Big) & \text{and} & y_{p}(t) &= y \Big( t, x_{0}, y_{0}, t_{0} \Big) \\ x_{st} \Big( t_{0} \Big) &= x \Big( t, x_{0}, y_{0}, t_{0} \Big) & \text{and} & y_{st} \Big( t_{0} \Big) &= y \Big( t, x_{0}, y_{0}, t_{0} \Big) \end{aligned}$$

which gives the streakline at t, where  $x_0$ ,  $y_0$  is the point at which dye is released ( $t_0$  is varied from 0 to t)

### Assumption: 2D flow

For pathlines	$u_p = \frac{dx}{dt} = B \cdot x \cdot (1 + A \cdot t)$ $A = 0.5 \frac{1}{s}$	$B = 1  \frac{1}{s}  v_p = \frac{dy}{dt} = C \cdot y \qquad C = 1  \frac{1}{s}$
So, separating variables	$\frac{\mathrm{d}x}{\mathrm{x}} = \mathrm{B} \cdot (1 + \mathrm{A} \cdot \mathrm{t}) \cdot \mathrm{d}\mathrm{t}$	$\frac{\mathrm{d}y}{\mathrm{y}} = \mathrm{C} \cdot \mathrm{d}t$
Integrating	$\ln\left(\frac{x}{x_0}\right) = B \cdot \left(t - t_0 + A \cdot \frac{t^2 - t_0^2}{2}\right)$	$\ln\left(\frac{y}{y_0}\right) = C \cdot \left(t - t_0\right)$
	$x = x_0 \cdot e^{B \cdot \left(t - t_0 + A \cdot \frac{t^2 - t_0^2}{2}\right)}$	$\mathbf{y} = \mathbf{y}_0 \cdot \mathbf{e}^{\mathbf{C} \cdot \left(t - t_0\right)}$
The pathlines are	$x_{p}(t) = x_{0} \cdot e^{B \cdot \left(t - t_{0} + A \cdot \frac{t^{2} - t_{0}^{2}}{2}\right)}$	$y_{p}(t) = y_{0} \cdot e^{C \cdot (t-t_{0})}$

where  $x_0$ ,  $y_0$  is the position of the particle at  $t = t_0$ . Re-interpreting the results as streaklines:

$$x_{st}(t_0) = x_0 \cdot e^{B \cdot \left(t - t_0 + A \cdot \frac{t^2 - t_0^2}{2}\right)} \qquad \qquad y_{st}(t_0) = y_0 \cdot e^{C \cdot (t - t_0)}$$

The streaklines are then

where  $x_0$ ,  $y_0$  is the point at which dye is released ( $t_0$  is varied from 0 to t)

For streamlines  $\frac{v}{u} = \frac{dy}{dx} = \frac{C \cdot y}{B \cdot x \cdot (1 + A \cdot t)}$ So, separating variables  $(1 + A \cdot t) \cdot \frac{dy}{y} = \frac{C}{B} \cdot \frac{dx}{x}$  which we can integrate for any given t (t is treated as a constant) Integrating  $(1 + A \cdot t) \cdot \ln(y) = \frac{C}{B} \cdot \ln(x) + \text{const}$ The solution is  $y^{1+A \cdot t} = \text{const} \cdot x^{\frac{C}{B}}$ For particles at (1,1) at t = 0, 1, and 2s y = x  $y = x^{\frac{2}{3}}$   $y = x^{\frac{1}{2}}$ 



2.29 Streaklines are traced out by neutrally buoyant marker fluid injected into a flow field from a fixed point in space. A particle of the marker fluid that is at point (x, y) at time t must have passed through the injection point  $(x_0, y_0)$  at some earlier instant  $t = \tau$ . The time history of a marker particle may be found by solving the pathline equations for the initial conditions that  $x = x_0$ ,  $y = y_0$  when  $t = \tau$ . The present locations of particles on the streakline are obtained by setting  $\tau$ equal to values in the range  $0 \le \tau \le t$ . Consider the flow field  $\vec{V} = ax(1+bt)\hat{t}+cy\hat{j}$ , where a=c=1 s<sup>-1</sup> and b=0.2 s<sup>-1</sup>. Coordinates are measured in meters. Plot the streakline that passes through the initial point  $(x_0, y_0) = (1, 1)$ , during the interval from t=0 to t=3 s. Compare with the streamline plotted through the same point at the instants t = 0, 1, and2 s.

#### Given: Velocity field

Find: Plot of streakline for t = 0 to 3 s at point (1,1); compare to streamlines through same point at the instants t = 0, 1and 2 s

Solution:

Governing equations:	For pathlines	$u_p = \frac{dx}{dt}$	$v_p = \frac{dy}{dt}$	For streamlines	$\frac{v}{u} = \frac{dy}{dx}$
----------------------	---------------	-----------------------	-----------------------	-----------------	-------------------------------

Following the discussion leading up to Eq. 2.10, we first find equations for the pathlines in form

$$\begin{aligned} x_{p}(t) &= x \Big( t, x_{0}, y_{0}, t_{0} \Big) & \text{and} & y_{p}(t) &= y \Big( t, x_{0}, y_{0}, t_{0} \Big) \\ x_{st} \Big( t_{0} \Big) &= x \Big( t, x_{0}, y_{0}, t_{0} \Big) & \text{and} & y_{st} \Big( t_{0} \Big) &= y \Big( t, x_{0}, y_{0}, t_{0} \Big) \end{aligned}$$

which gives the streakline at t, where  $x_0$ ,  $y_0$  is the point at which dye is released ( $t_0$  is varied from 0 to t)

Assumption: 2D flow

For pathlines

Integrating

For pathlines  

$$u_{p} = \frac{dx}{dt} = a \cdot x \cdot (1 + b \cdot t) \qquad a = 1 \quad \frac{1}{s} \qquad b = \frac{1}{5} \quad \frac{1}{s} \quad v_{p} = \frac{dy}{dt} = c \cdot y \qquad c = 1 \quad \frac{1}{s}$$
So, separating variables  

$$\frac{dx}{x} = a \cdot (1 + b \cdot t) \cdot dt \qquad \qquad \frac{dy}{y} = c \cdot dt$$
Integrating  

$$\ln\left(\frac{x}{x_{0}}\right) = a \cdot \left(t - t_{0} + b \cdot \frac{t^{2} - t_{0}^{2}}{2}\right) \qquad \qquad \ln\left(\frac{y}{y_{0}}\right) = c \cdot (t - t_{0})$$

$$x = x_0 \cdot e^{a \cdot \left(t - t_0 + b \cdot \frac{t^2 - t_0^2}{2}\right)} \qquad \qquad y = y_0 \cdot e^{c \cdot \left(t - t_0\right)}$$

$$x_{p}(t) = x_{0} \cdot e^{a \cdot \left(t - t_{0} + b \cdot \frac{t^{2} - t_{0}^{2}}{2}\right)}$$

$$y_{p}(t) = y_{0} \cdot e^{c \cdot \left(t - t_{0}\right)}$$

The pathlines are

where  $x_0$ ,  $y_0$  is the position of the particle at  $t = t_0$ . Re-interpreting the results as streaklines:

The streak lines are then 
$$x_{st}(t_0) = x_0 \cdot e^{-a \cdot \left(t - t_0 + b \cdot \frac{t^2 - t_0^2}{2}\right)}$$
  $y_{st}(t_0) = y_0 \cdot e^{-c \cdot (t - t_0)}$   
where  $x_0, y_0$  is the point at which dye is released ( $t_0$  is varied from 0 to t)  
For streamlines  $\frac{v}{u} = \frac{dy}{dx} = \frac{c \cdot y}{a \cdot x \cdot (1 + b \cdot t)}$   
So, separating variables  $(1 + b \cdot t) \cdot \frac{dy}{y} = \frac{c}{a} \cdot \frac{dx}{x}$  which we can integrate for any given t (t is treated as a constant)  
Integrating  $(1 + b \cdot t) \cdot \ln(y) = \frac{c}{a} \cdot \ln(x) + \text{const}$   
The solution is  $y^{1+b \cdot t} = \text{const} \cdot x^{\frac{c}{a}}$   
For particles at (1,1) at t = 0, 1, and 2s  $y = x$   $y = x^{\frac{2}{3}}$   $y = x^{\frac{1}{2}}$ 





x (m)

**2.30** Consider the flow field  $\vec{V} = axt\hat{i} + b\hat{j}$ , where  $a = 1/4 \text{ s}^{-2}$  and b = 1/3 m/s. Coordinates are measured in meters. For the particle that passes through the point (x, y) = (1, 2) at the instant t = 0, plot the pathline during the time interval from t = 0 to 3 s. Compare this pathline with the streakline through the same point at the instant t = 3 s.

### **Given:** Velocity field

**Find:** Plot of pathline for t = 0 to 3 s for particle that started at point (1,2) at t = 0; compare to streakline through same point at the instant t = 3

### Solution:

Governing equations:	For pathlines	$u_p = \frac{dx}{dt}$	$v_p = \frac{dy}{dt}$
		ui	uı

Following the discussion leading up to Eq. 2.10, we first find equations for the pathlines in form

$$x_{p}(t) = x(t, x_{0}, y_{0}, t_{0})$$
 and  $y_{p}(t) = y(t, x_{0}, y_{0}, t_{0})$ 

$$\mathbf{x}_{st}(t_0) = \mathbf{x}(t, \mathbf{x}_0, \mathbf{y}_0, t_0) \qquad \text{and} \qquad \mathbf{y}_{st}(t_0) = \mathbf{y}(t, \mathbf{x}_0, \mathbf{y}_0, t_0)$$

which gives the streakline at t, where  $x_0$ ,  $y_0$  is the point at which dye is released ( $t_0$  is varied from 0 to t)

### Assumption: 2D flow

For pathlines	$u_p = \frac{dx}{dt} = a \cdot x \cdot t$	$a = \frac{1}{4}  \frac{1}{\frac{2}{s}}$	$b = \frac{1}{3}  \frac{m}{s}$	$v_p = \frac{dy}{dt} = b$
So, separating variables	$\frac{\mathrm{d}x}{\mathrm{x}} = \mathrm{a} \cdot \mathrm{t} \cdot \mathrm{d}\mathrm{t}$			$dy = b \cdot dt$
Integrating	$\ln\left(\frac{x}{x_0}\right) = \frac{a}{2} \cdot \left(t^2 - t_0^2\right)$			$y - y_0 = b \cdot (t - t_0)$
	$\mathbf{x} = \mathbf{x}_0 \cdot \mathbf{e}^{\frac{\mathbf{a}}{2} \cdot \left(t^2 - t_0^2\right)}$			$y = y_0 + b \cdot (t - t_0)$
The pathlines are	$x_{p}(t) = x_{0} \cdot e^{\frac{a}{2} \cdot \left(t^{2} - t_{0}^{2}\right)}$			$y_{p}(t) = y_{0} + b \cdot (t - t_{0})$

where  $x_0$ ,  $y_0$  is the position of the particle at  $t = t_0$ . Re-interpreting the results as streaklines:

 $x_{st}(t_0) = x_0 \cdot e^{\frac{a}{2} \cdot (t^2 - t_0^2)} \qquad y_{st}(t_0) = y_0 + b \cdot (t - t_0)$ 

The pathlines are then

where  $x_0$ ,  $y_0$  is the point at which dye is released ( $t_0$  is varied from 0 to t)



**2.31** A flow is described by velocity field  $\vec{V} = ay^2\hat{i} + b\hat{j}$ , where  $a = 1 \text{ m}^{-1}\text{s}^{-1}$  and b = 2 m/s. Coordinates are measured in meters. Obtain the equation for the streamline passing through point (6, 6). At t = 1 s, what are the coordinates of the particle that passed through point (1, 4) at t = 0? At t = 3 s, what are the coordinates of the particle that passed through point (-3, 0) 2 s earlier? Show that pathlines, streamlines, and streaklines for this flow coincide.

**Given:** 2D velocity field

**Find:** Streamlines passing through (6,6); Coordinates of particle starting at (1,4); that pathlines, streamlines and streaklines coincide

### Solution:

For streamlines	$\frac{v}{u} = \frac{dy}{dx} = \frac{b}{a \cdot y^2}$ or	$\int a \cdot y^2  dy = \int b  dx$	
Integrating	$\frac{\mathbf{a} \cdot \mathbf{y}^3}{3} = \mathbf{b} \cdot \mathbf{x} + \mathbf{c}$		
For the streamline through point (6,6)	c = 60 and	$y^3 = 6 \cdot x + 180$	
For particle that passed through $(1,4)$ at $t = 0$	$u = \frac{dx}{dt} = a \cdot y^2$	$\int 1  dx = x - x_0 = \int$	$a \cdot y^2 dt$ We need $y(t)$
	$v = \frac{dy}{dt} = b$	$\int 1  dy = \int b  dt$	$\mathbf{y} = \mathbf{y}_0 + \mathbf{b} \cdot \mathbf{t} = \mathbf{y}_0 + 2 \cdot \mathbf{t}$
Then	$x - x_0 = \int_0^t a \cdot (y_0 + b \cdot t)^2 dt$	$\mathbf{x} = \mathbf{x}_0 + \mathbf{a} \cdot \left( \mathbf{y}_0^2 \cdot \mathbf{t} + \mathbf{b} \cdot \mathbf{y}_0^2 \right)$	$y_0 \cdot t^2 + \frac{b^2 \cdot t^3}{3}$
Hence, with $x_0 = 1$ $y_0 = 4$	$x = 1 + 16 \cdot t + 8 \cdot t^2 + \frac{4}{3} \cdot t^3$	At $t = 1 s$	x = 26.3·m
	$y = 4 + 2 \cdot t$		$y = 6 \cdot m$
For particle that passed through $(-3,0)$ at t = 1	$\int 1  dy = \int b  dt$	$y = y_0 + b \cdot \left( t - t_0 \right)$	
$x - x_0 = \int_{t_0}^t a \cdot (y_0 + b \cdot t)^2 dt$	$\mathbf{x} = \mathbf{x}_0 + \mathbf{a} \cdot \left[ \mathbf{y}_0^2 \cdot \left( \mathbf{t} - \mathbf{t}_0 \right) + \mathbf{b} \cdot \mathbf{y} \right]$	$t_0 \cdot (t^2 - t_0^2) + \frac{b^2}{3} \cdot (t^3 - t_0^2) + \frac{b^2}$	$t_0^3$
Hence, with $x_0 = -3$ , $y_0 = 0$ at $t_0 = 1$	$x = -3 + \frac{4}{3} \cdot (t^3 - 1) = \frac{1}{3} \cdot (4 \cdot t^3)$	$y = 2 \cdot (t - 13)$	1)
Evaluating at t = 3	$x = 31.7 \cdot m$	$y = 4 \cdot m$	

This is a steady flow, so pathlines, streamlines and streaklines always coincide

# Fox and McDonalds Introduction to Fluid Mechanics 8th Edition Pritchard Solutions Manual

Full Download: http://alibabadownload.com/product/fox-and-mcdonalds-introduction-to-fluid-mechanics-8th-edition-pritchard-sol

Problem 2.32

[Difficulty: 3]

S

2.32 Tiny hydrogen bubbles are being used as tracers to visualize a flow. All the bubbles are generated at the origin (x = 0, y = 0). The velocity field is unsteady and obeys the equations:
u = 1 m/s v = 2 m/s 0 ≤ t < 2 s u = 0 v = -1 m/s 0 ≤ t ≤ 4 s</li>
Plot the pathlines of bubbles that leave the origin at t = 0, 1, 2, 3, and 4 s. Mark the locations of these five bubbles at t = 4 s.

### Solution

The particle starting at t = 3 s follows the particle starting at t = 2 s; The particle starting at t = 4 s doesn't move!

Pathlines:	Startin	g at $t = 0$ Starting at $t = 1$ s		Starting at $t = 2 s$		s	Streakline at t = 4				
t	х	У	x	У		x	У		x	У	
0.00	0.00	0.00							2.00	2.00	
0.20	0.20	0.40							1.80	1.60	
0.40	0.40	0.80							1.60	1.20	
0.60	0.60	1.20							1.40	0.80	
0.80	0.80	1.60							1.20	0.40	
1.00	1.00	2.00	0.00	0.00					1.00	0.00	
1.20	1.20	2.40	0.20	0.40					0.80	-0.40	
1.40	1.40	2.80	0.40	0.80					0.60	-0.80	
1.60	1.60	3.20	0.60	1.20					0.40	-1.20	
1.80	1.80	3.60	0.80	1.60					0.20	-1.60	
2.00	2.00	4.00	1.00	2.00		0.00	0.00		0.00	-2.00	
2.20	2.00	3.80	1.00	1.80		0.00	-0.20		0.00	-1.80	
2.40	2.00	3.60	1.00	1.60		0.00	-0.40		0.00	-1.60	
2.60	2.00	3.40	1.00	1.40		0.00	-0.60		0.00	-1.40	
2.80	2.00	3.20	1.00	1.20		0.00	-0.80		0.00	-1.20	
3.00	2.00	3.00	1.00	1.00		0.00	-1.00		0.00	-1.00	
3.20	2.00	2.80	1.00	0.80		0.00	-1.20		0.00	-0.80	
3.40	2.00	2.60	1.00	0.60		0.00	-1.40		0.00	-0.60	
3.60	2.00	2.40	1.00	0.40		0.00	-1.60		0.00	-0.40	
3.80	2.00	2.20	1.00	0.20		0.00	-1.80		0.00	-0.20	
4.00	2.00	2.00	1.00	0.00		0.00	-2.00		0.00	0.00	

